



UNIVERSIDAD PERUANA
CAYETANO HEREDIA
ESCUELA DE POSGRADO

NEEDS, ACCEPTABILITY, AND VALUE OF
HUMANITARIAN MEDICAL ASSISTANCE IN
REMOTE PERUVIAN AMAZON RIVERINE
COMMUNITIES

TESIS PARA OPTAR EL GRADO DE MAESTRO EN
CIENCIAS EN INVESTIGACIÓN EPIDEMIOLÓGICA

JUAN FRANCISCO SANCHEZ CHIROQUE

LIMA - PERÚ

2016

Asesor de Tesis

Andres Guillermo Lescano Guevara

Acknowledgments

We thank all of the study participants and all of the fieldworkers for collecting the information for this article, especially our supervisors, Connie Fernandez and Edson Maguiña. In addition, we would like to thank Kimberly A. Edgel for critical review of this manuscript and Hugo Valdivia for his help with mapping. Also, we thank microscopists in the communities, Lima, and Iquitos for their outstanding diagnostic assistance. We are thankful for the support and guidance received from program faculty and fellow students.

Financial support

This study was supported by the U.S. Southern Command (SOUTHCOM) under work unit 847705 82000 25GB B0016 and the National Institutes of Health/Fogarty International Center (NIH/FIC) under Training Grant 2D43 TW007393

RESUMEN

Existe un gran debate acerca de la necesidad, la aceptabilidad y el valor de la asistencia médica humanitaria. Realizamos un estudio transversal en 457 niños menores de 5 años de cuatro comunidades ribereñas remotas de la Amazonía peruana y recogimos las medidas antropométricas, muestras de sangre (1-4 años), y muestras de heces. Grupos focales y entrevistas a informantes clave evaluaron las perspectivas respecto a la ayuda médica brindada por extranjeros. La prevalencia de la desnutrición crónica, anemia y parásitos intestinales fueron 20%, 37% y 62%, respectivamente. La infección con múltiples parásitos, usualmente geohelminthos, se detectó en el 41% de los niños. La prevalencia de parásitos intestinales tanto individuales como poliparasitismo aumentaba con la edad. Los participantes de las comunidades más pequeñas y menos expuestas a los extranjeros expresaron falta de confianza y miedo a ellos. Sin embargo, los participantes de todas las comunidades tuvieron positivos comentarios acerca de los extranjeros que acuden para proporcionar apoyo en salud. Las prevalentes necesidades de salud, como las infecciones parasitarias y anemia, pueden ser abordadas por las intervenciones médicas de corto plazo. Hay una adecuada actitud receptiva y aceptación de la asistencia médica entregada por personal extranjero.

ABSTRACT

Much debate exists regarding the need, acceptability, and value of humanitarian medical assistance. We conducted a cross-sectional study on 457 children under 5 years from four remote riverine communities in the Peruvian Amazon and collected anthropometric measures, blood samples (1–4 years), and stool samples. Focus groups and key informant interviews assessed perspectives regarding medical aid delivered by foreigners. The prevalence of stunting, anemia, and intestinal parasites was 20%, 37%, and 62%, respectively. Infection with multiple parasites, usually geohelminths, was detected in 41% of children. The prevalence of intestinal parasites both individual and polyparasitism increased with age. Participants from smaller communities less exposed to foreigners expressed lack of trust and fear of them. However, participants from all communities were positive about foreigners visiting to provide health support. Prevalent health needs such as parasitic infections and anemia may be addressed by short-term medical interventions. There is a perceived openness to and acceptability of medical assistance delivered by foreign personnel.

INDICE

CONTENIDO	Páginas
I. Introduction	01
II. Materials and Methods	04
III. Results	12
IV. Discussion	18
V. References	23

I. INTRODUCTION

Resource-limited areas often have limited financial capital, human resources, and infrastructure in a wide array of areas such as education, security, transport, and health (1-3). The Department of Loreto is located in the Peruvian Amazon basin, the region with the worst health indicators (4) and the highest risk of infectious diseases (5) in Peru. This region also has the highest incidence of malaria, diarrheal disease, and dengue in the country, similar to other tropical rural areas in Latin America and the Caribbean (6).

The people of the Amazon region often live in small villages along the Amazon River and its tributaries (5), some of them in very remote river basins. Most of this population consider themselves as “mestizo,” descendants from indigenous populations who use and accept occidental culture (7). Poor health status across all of these populations, in addition to inadequate health infrastructure and resources, limit their access to quality health services (5). Young children in the region, particularly those less than 5 years of age, frequently experience nutritional deficiencies, with increased risk of resulting developmental disorders at critical stages of growth (8). The remote location of many of these populations and large distances between communities creates further challenges (7). Therefore, the Peruvian Ministry of Health (MoH) and other governmental agencies face significant difficulties when trying to provide necessary health services.

In order to provide medical care and improve the well-being of Amazonian and other vulnerable populations worldwide, non-governmental organizations (NGOs), religious organizations, and foreign institutions carry out humanitarian assistance (HA). HA programs aim to address health disparities through provision of short-term basic healthcare services and occasionally by supporting infrastructure development to meet basic human needs such as water and sewage (9).

Health-wise, HA programs focus mainly on acute conditions that can be addressed with a single intervention, such as treatment of intestinal parasites, fungal skin diseases, and nutritional deficiencies (10). These programs are generally carried out by trained health personnel from different specialties (pediatrics, nutrition, dentistry, dermatology) and often focus on the most vulnerable populations such as children and pregnant women (11). The Department of Defense of the United States of America (US-DoD), often conducts HA activities in developing countries as part of “Medical Readiness Training Exercise” (MEDRETEs). The main goal of MEDRETEs is to train active duty personnel of the US Armed Forces in the delivery of medical care in remote areas with limited access to healthcare. Secondary benefits of MEDRETEs include provision of humanitarian and civic assistance, healthcare services and treatment (12, 13).

HA activities are often criticized for poor integration with local health services and limited evidence of a lasting positive impact, mainly because of their short duration and lack of intensity and continuity (9, 14). In an attempt to address these criticisms, the U.S. Air Force selected four remote riverine communities in the Amazon rainforest for repeated MEDRETE interventions in consecutive years starting in 2009. The expectation was that through repeated interventions in those communities, HA could potentially continue to improve the health of community members, especially children. Attempts were also made to improve the sustainability of the impact of interventions by providing training to local MoH personnel stationed in the area (15, 16), while accomplishing the primary training goal of MEDRETEs.

The possibility of offering repeated health interventions in these communities provided a unique opportunity to assess the impact of the MEDRETE HA activities. We performed a baseline evaluation of the health status of children under 5 years in these four communities before the beginning of HAs because children in this age range are one of the main target populations for

their activities. The main goals of this study were: 1) to identify health needs in vulnerable population amenable to be addressed by HA and suggest potential interventions, 2) assess the acceptability of HA, and 3) to provide baseline information about health status of children that could be used to evaluate the impact of the HA actions.

II. MATERIALS AND METHODS

Settings

Between January to March 2009, we conducted a cross-sectional health assessment prior to a planned HA intervention. We included four remote riverine communities in the western region of the Department of Loreto, located in the Peruvian Amazon Basin: Saramiriza (population: 1947) and Puerto America (population: 601), along the Marañón River; and Lagunas (population: 6317) and Santa Cruz (population: 1910), along the Huallaga River (Figure 1). Public services in these communities are very limited (Table 1). The weather in this area is rainy and tropical with an average relative humidity of 84% and a mean temperature of 27°C (22°C – 36°C). Rainfall increases between December and March (5). Travel time to the nearest large city, (Yurimaguas, population: 45756), is 2 to 12 hours by speed boat, and much longer via the slower boats commonly used for local transport. All four communities mainly have non-potable drinking water sources, poor sanitation facilities and practice open defecation, according to the categories reported in the JMP method by the WHO (17). Also, they have three to eight hours of electricity daily, depending of the community. River water is used by the population for cooking, drinking, and other daily activities, and the people from these communities eliminate litter and feces in the river or in the open field. A few members of the communities have access to latrines, but the latrine waste goes directly to the river.

Local health services are similarly limited. Most healthcare providers are recent graduates from different health professions (e.g., physicians, nurses, veterinarians) performing a year of rural service as part of the national Rural and Marginal Urban Areas Health Service (SERUMS) program. Saramiriza and Lagunas have the best-staffed health facilities; however, only two physicians and two nurses provided services at each facility during the time of the study. The

health facility in Puerto America is the smallest (80 m²) and had no physicians, but only one professional midwife, one nurse, and two technicians on staff. Santa Cruz has a health facility with better infrastructure than Puerto America but both only had a physician and nurse. Also, no health facility possessed laboratory equipment nor a well-stocked pharmacy. The nearest hospital with capacity to address more complex medical problems is located in Yurimaguas (Table 1).

Fieldwork

We performed a population census to identify eligible individuals in the four communities. Then, we randomly selected prospective participants for a baseline assessment in three study groups: children under 5 years, 15-49 years “adult” caretakers of children under 5 years, and 15-49 years “adult” female and male sampled independently. The “adult” age range (15-49 years) was selected for comparability with the Demographic Health Surveys (DHS) standard age group used worldwide. The sample size of each study group was proportional to the number of inhabitants in that group for each community. Four 2-person teams conducted field operations for 52 days. Fieldworkers were health personnel and JFS and AMB trained them over three days in all study procedures (surveys and sample collection). A pilot study was conducted to confirm that all the procedures were understood. JFS was with the team throughout field activities and had daily meetings to address questions or doubts. The local population speaks Spanish and interpreters were not needed. In this manuscript, we only present results from children under 5 years, the most vulnerable of the three populations.

The data collected with qualitative (focus groups and key-personnel interviews) and quantitative (questionnaires and biological samples) components was used to evaluate the potential value of HA activities according to the community members and local MoH personnel.

Surveys

Trained personnel conducted interviews during household visits using questionnaires previously applied in Peru and piloted in the field by JFS. Two questionnaires were used: a census form and a survey regarding children's health. The census form was based on the 2007 Peruvian census questionnaire. Key data was recorded, including geographic coordinates, basic household characteristics, and socio-demographic information from each household member adjusting content for readability. The children questionnaire followed closely the standard DHS survey applied in Peru since 1986. The child survey was filled by a fieldworker with the information provided by the caretaker. The survey asked caretakers about one of their children under five years who lived in the house. Caretakers were asked about their threshold for seeking healthcare for their child and their child's perceived health status, immunizations received, and recent illnesses. Additionally, fieldworkers requested the child's immunization card to collect vaccination records. If unavailable, the caretaker asked about the immunizations received by the child. Efforts were made to apply the questionnaires in a private setting with an interviewer of the same sex as the caretaker. Also, the survey measured some benefits in members of the communities and local MoH personnel such as perception about MoH personnel and use of healthcare facilities. We did not evaluate the benefits in US Air Force personnel in training during this MEDRETE.

Anthropometrics

Height and weight were measured in all participants. Standard scales following UNICEF standards were used to measure weight (Seca 840, Hamburg, Germany). Stadiometers were locally made following specifications from UNICEF and the Peruvian National Center for Feeding and Nutrition and validated prior to the study. Weight and height were measured with a

precision of ± 100 g and ± 0.1 cm, respectively. Participants were asked to wear the minimal clothing needed and no shoes to avoid overestimations, and the weight of the clothes worn was discounted using standard cloth weight tables. The recumbent length of children under 2 years was measured lying on their backs and the height of children 2 years and older was measured standing up. Two members of the field team worked together to carry out the anthropometric measures. Stunting and wasting were defined as height-for-age and weight-for-height less than -2 z-scores below the median of the WHO child growth standards, respectively.

Assessment of malaria and anemia

Peripheral blood obtained by finger prick was used to diagnose malaria and assess anemia by puncturing the third finger of the left hand of all participants age 1 year and over. Standard methods and universal precautions were used. Three or four drops were collected per child: two were used for malaria diagnosis by thick and thin blood smears stained on-site with Giemsa stain and one or two for anemia testing. Smears were examined for malaria in batch at the local health facilities and confirmed by a second reading at the U.S. Naval Medical Research Unit No. 6 (NAMRU-6) in Iquitos (18). The result of the first reading was reported to parents within 24 hours of sample collection following MoH reporting practices. One or two drops were placed in a microcuvette for on-site hemoglobin testing using a Hemocue® electronic measurement device (HemoCue HB 201+ Analyzer, HemoCue AB, Angelholm, Sweden). Upon obtaining the result, the fieldworker reported to the caretaker whether or not the child had anemia according to international standards (Hemoglobin < 11 g/dL in children). Results were also recorded on the questionnaire.

Enteric parasites

We collected one stool sample from each participant. Twenty grams of stool were placed into sodium acetate–acetic acid–formalin solution to preserve intestinal parasites. Specimens were also placed in cryovials and kept in a nitrogen tank to preserve fresh stool. The samples were processed with the Ritchie concentration technique (19) using ethyl acetate instead of ether. Stool smears were read by experienced microscopists under direct observation using saline and Lugol's iodine solution at NAMRU-6 in Lima and Iquitos to identify intestinal parasites. Unconcentrated stool was spotted onto slides for modified Kinyoun's stain to look for *Cryptosporidium* spp. and *Cyclospora cayetanensis*.

The cryovials were sent to the US Centers for Disease Control and Prevention (CDC) Division of Parasitic Diseases and Malaria in Atlanta, Georgia, for molecular testing for *Entamoeba histolytica/Entamoeba dispar*, *Giardia lamblia*, and *Cryptosporidium* spp. Four hundred μ L of fecal suspension were processed as instructed with the FastDNA SPIN kit for Soil (MP Biomedicals, Irvine, CA). For conventional polymerase chain reaction (PCR), we used a region of 877 base pairs (bp) of the small subunit ribosomal RNA (SSU rRNA) and for real-time PCR (RT-PCR), we used TaqMan RT-PCR based on the SSU rRNA gene. Primers Ehd-239F and Ehd-88R amplified a 172 bp fragment of the SSU rRNA gene of *E. histolytica/E. dispar*. The amoebae were discriminated with species-specific probes *E. histolytica*-96T and *E. dispar*-96T, respectively. For *Giardia lamblia*, specimens were confirmed by TaqMan RT-PCR targeting a 105 bp fragment of the SSU rRNA gene. The molecular characterization of *Giardia lamblia* was done using nested PCR to amplify a ~532 bp fragment of the triose phosphate isomerase (TPI) gene. The genotyping of *Cryptosporidium* spp. used a SSU rRNA–based PCR-restriction fragment length polymorphism that differentiated *Cryptosporidium* spp. Identification of selected

genotypes was confirmed by DNA sequence analysis of these amplicons. The subtypes of *Cryptosporidium hominis* were determined by sequence analysis of a ~850 bp fragment of the 60-kDa glycoprotein gene (20).

Qualitative assessment

The objective of the qualitative component was to explore local perspectives regarding health and well-being and factors that influence health and humanitarian aid from potential direct and indirect beneficiaries. Seventy individuals were recruited using purposive sampling. Four focus groups were conducted, one in each community, with a total of 34 participants overall and 8-10 participants per focus group. Additionally, 36 in-depth, semi-structured interviews of indirect beneficiaries were conducted with formal and informal community leaders such as healthcare personnel, mayors, heads of women's organizations, heads of community kitchens (*comedores populares*), and the Glass of Milk program. We selected community leaders for interviews because they have important opinions regarding issues of interest to the community since that is part of their role as leaders. Also, they tend to be outspoken and their opinion is respected and rarely challenged by community members; for this reason, is not productive to include them on focus groups. Focus group participants and interviewees were recruited by the qualitative research assistant in cooperation with local collaborators in each community. Both focus groups and interviews addressed diverse topics. First, they engaged participants about how they define and perceive health and well-being and the factors that contribute to each and about the current state of health and well-being among residents of their community. Then, they asked participants about common health problems among children. Next, they asked participants about factors and resources that both improve and weaken their health and well-being. Finally, the focus groups and interviews asked participants about their perceptions regarding the presence of foreigners

and HA in their communities. The focus groups and interviews were carried out by the qualitative research assistant, a Peruvian sociologist with extensive training in qualitative research and ethics and extensive experience implementing qualitative data collection with diverse populations in Peru. The research assistant and AMB, a qualitative researcher, analyzed the data using a grounded theory approach. Following thorough reading of the focus groups and interviews, they developed an initial codebook. Then, they separately coded one focus group and two interview transcripts and then jointly reviewed the coding to standardize the coding process and refine the codebook. Next, the research assistant coded the remaining transcripts. Finally, the research assistant and AMB worked together to synthesize the results, comparing similarities and differences across the four communities.

The audio of the narrative portion of the focus groups and all the in-depth interviews were recorded with the consent of the participants. The recordings and notes were transcribed and the resulting transcripts were analyzed using a grounded-theory approach. Once each set of interviews was complete, coding was carried out using the Atlas-ti software program (Scientific Software Development GmbH, 2005). The participatory portion of the focus groups was recorded jointly with the participants, in addition to notes that were taken during and after the focus group by the facilitator. All recorded material was then analyzed by an experienced PhD level internationally trained scientist (AMB) who resides in Peru for eight years, drawing parallels and differences across the study communities.

Statistics

Dichotomous outcomes such as disease prevalence are presented as proportions and numerical outcomes are presented as means or medians depending on the presence of outliers. We calculated 95% confidence intervals with exact binomial methods for the main dichotomous

outcomes. Comparison of proportions across years used Chi2 and Fisher's exact tests if needed, while we used Student's T and Mann-Whitney Wilcoxon tests for numerical outcomes depending on the presence of influential outliers. All analysis were conducted with Stata 13.0 (Stata Corp., College Station, TX) using a 0.05 significance level.

Ethics

The Institutional Review Board (IRB) of NAMRU-6 approved the study protocol in compliance with all applicable US federal regulations governing the protection of human subjects. In addition, the Loreto Regional Health Directorate (DIRESA) in Iquitos and the Alto Amazonas Health Network in Yurimaguas approved the study on behalf of the Peruvian MoH. Work at CDC was approved by the CDC-IRB. Additionally, community consent was obtained from community leaders identified by local MoH personnel, as a collaborative effort between study personnel and local MoH personnel. Each participant in the quantitative and qualitative components as well as the parents of all children engaged in a standardized informed consent process and provided signed informed consent prior to initiating participation. All approvals were obtained before starting any activities in each community.

III. RESULTS

Six communities were pre-selected in coordination with the local Peruvian MoH according to criteria such as lack of basic services (sanitation, safe water, and electricity), access by large ships from the Peruvian Navy, and presence of health services. The team visited all the communities and four were chosen. We evaluated 1752 households (65% of the total) in the four communities during the census and enrolled 457 children under 5 years (39% of the total) and their caretakers.

Sociodemographic characteristics

The study sample was approximately distributed proportional to the size of each community, and 58% of the children were from Lagunas. The number of males and females was similar (49% vs. 51%). Participants' mean age (\pm standard deviation) was 29 ± 18.5 months and there was no difference in average age between communities ($p=0.096$). The demographic characteristics of the children that participated are presented in Table 2.

Nutritional status

As shown in Table 3, the prevalence of stunting in all children was 20% (81/406). Puerto America (33%) had the highest prevalence, but there were no statistical differences between the communities ($p=0.466$). The highest prevalence of stunting occurred in children over 1 year ($p<0.001$). The overall prevalence of wasting was 3% (13/406), very similar among the communities ($p=0.929$) but lower in children who were 3-4 years in comparison with children 1-2 years (2% vs. 8%, $p=0.022$).

The mean hemoglobin level in children aged 1-4 years was 11.2 ± 1.1 mg/dl (range: 7.3 – 14.6) and the prevalence of anemia was 37%. Hemoglobin levels were different across communities ($p=0.003$), with the highest mean in Saramiriza (11.5 mg/dl) and the lowest in Lagunas (11.1

mg/dl); however, this did not result in significant differences ($\Delta \leq 17\%$) in anemia across the four towns ($p=0.117$). We also found lower hemoglobin levels in children 1-2 years compared to children 3-4 years (10.8 versus 11.6 mg/dl, $p<0.001$) and corresponding higher levels of anemia in 1-2 years versus 3-4 years (55% versus 23%, $p<0.001$).

Intestinal parasites and malaria

The prevalence of intestinal parasites was 62% (Table 4). The most common were *Ascaris lumbricoides* (29%), *Giardia lamblia* (20%), *Trichuris trichiura* (16%), hookworms (5%), and *Strongyloides stercoralis* (2%). The four communities presented statistically different prevalences of the four parasites, potentially due to differences in infrastructure (urbanization) and access to antiparasitic drugs between communities: *Giardia lamblia* ($p=0.011$), *Trichuris trichiura* ($p<0.001$), *Entamoeba histolytica/dispar* ($p=0.001$) and *Strongyloides stercoralis* ($p=0.013$). The prevalence of most intestinal parasites increased with age. Polyparasitism was similarly frequent across the four communities ($p=0.179$), but its prevalence also increased with age ($p<0.001$).

Thirty-four samples positive by microscopy for *E. histolytica/dispar* (8%) were evaluated by conventional PCR and RT-PCR. Eleven samples (32%) positive for *E. dispar* by conventional PCR and 18 (53%) were positive by RT-PCR. No sample was positive for *E. histolytica* by conventional PCR or RT-PCR. Seventy-two of 80 samples (90%) positive by microscopy for *G. lamblia* were confirmed positive by RT-PCR. Forty-three of these specimens were successfully amplified by TPI PCR and sequenced. Three children (7%) had assemblage A, and 40 (93%) had assemblage B. BS1 was the most frequent sub-assemblage, found in 35% of the samples (15/43). Six participants were simultaneously infected with two B sub-assemblages.

[Coccidian protozoans](#) were also found by modified acid fast staining. Twenty samples (4%) were identified as *Cryptosporidium* spp.; also three were reported as *Cyclospora cayetanensis*. Seven samples identified as *Cryptosporidium* spp. were successfully characterized at the species level. The species detected were *Cryptosporidium hominis* (n=5) and *C. canis* (n=2). Subtyping was successful in four of five participants with *C. hominis*: subtypes IaA13R7 (n=1), IbA10G2 (n=2), and IaA11G3T3 (n=1). None of the *C. cayetanensis* were confirmed by molecular methods.

Direct microscopic evaluation of 327 blood smears revealed no evidence of *Plasmodium* or filariasis infection in any participant. Quality control was performed for expert microscopists in NAMRU-6, Iquitos and a concordance of 100% was found with the results from local MoH microscopists.

Immunizations and reported health status

We assessed immunizations in children over 1 year-old using the information collected from the vaccine cards (41%) or from caretakers (59%), if vaccine cards were not available. We found that only 71% of children over 1 year completed the full Peruvian immunization schedule, which at that time included hepatitis B, polio, measles, mumps, *Haemophilus influenzae* type b, and yellow fever (21), among other vaccines (Table 5).

Additionally, we evaluate children's health status asking caretakers if children had fever, cough or diarrhea in the two weeks before the survey and what they did to face this situation. They reported that 49% of their children had a cough, 34% had a fever, and 29% had diarrhea. When the children had fever, cough, or diarrhea, 72%, 60%, and 35% of the caretakers, respectively, sought care at an MoH health facility. Over half (57%) of caretakers whose children had a fever sought care during the first day with fever, in comparison to 37% of those with cough ($p<0.001$) and 26% of those with diarrhea ($p<0.001$). According to the caretakers, about half of the children

with fever (55%) or cough (45%) were seen by a physician compared to a quarter of diarrhea cases (25%). Nearly all children with fever in the last two weeks received antipyretics (89%) and only 10 children with fever (6%) received other drugs like antibiotics (amoxicillin) or corticosteroids. Half of the children (49%) received treatment for intestinal parasites in the last twelve months.

Finally, caretakers reported 61% of their children had good physical well-being and 67% had good emotional well-being. Fieldworkers also noted participants with decayed teeth and low visual acuity, another issue that can be addressed by short-term medical missions, although this was not systematically assessed in all participants.

Qualitative evaluations

Participants in focus groups and interviews defined health as the absence of illness and underscored that good health is important in order to be able to work. Well-being was described as a broad, comprehensive concept. Participants in most communities reported average health and well-being across adult and child populations, which leaders attributed to improvements in and expanded use of healthcare services. However, in the very small community of Puerto America, residents considered their health to be weak due to the lack of appropriate medical care and medications, as well as the numerous diseases faced by the population:

“The situation of the community and well-being is chaotic. [The population] doesn’t enjoy quality of life, also due to factors such as habits and lifestyles... The general population doesn’t have good habits, above all given what you see in terms of incidences of several illnesses, like ARIs, ADDs and parasitosis.” (Head of Health center, Male, Interview, Puerto América)

The child health ailments that participants considered to be most important were bronchial disease, diarrhea, and parasitic infection. Other problems perceived as common were: upper

respiratory infections in Saramiriza and Puerto America; malaria and other causes of fever in Saramiriza; anemia in Lagunas; and pneumonia and vomiting in Santa Cruz. Personal hygiene was mentioned as the factor that had the greatest positive influence on health in all of the communities, and the following factors were mentioned in three out of four communities: hand washing, washing and cooking food well, drinking boiled water, evaluation at MoH health facilities, and maintaining the cleanliness of the latrine. Lack of improved sanitation facilities was mentioned as the main factor that weakens or worsens health, in addition to lack of improved drinking water sources, lack of bathrooms or latrines, and littering. Participants also mentioned contamination by oil extraction companies and contamination of the river due to its use as a trash dumping area. All participants stated that most community members are informed about disease prevention practices, but that they still do not use them.

All four communities mentioned that they have three important resources in their communities: the MoH facility, the preschool, and schools in general. They spoke at length about pharmacies and traditional healers, both of which exist in greater number than skilled healthcare providers, although they also recognized that their local pharmacy personnel may not be trained healthcare providers. A healthcare professional interviewee mentioned that traditional healers tend to wait too long before referring their clients to formal health facilities.

In the larger community of Lagunas, participants reported more contact with foreigners, as the community is one of the entry points to an internationally recognized national reserve (Pacaya Samiria), and because they received health campaigns and visits by religious groups in the past. Participants were satisfied with these visits, especially since they felt they do not receive sufficient support from the MoH health system. The other communities, however, reported a minimal presence of foreigners. They said that non-Peruvians usually came to carry out research,

but the local inhabitants had not seen the results of the studies. All participants, and particularly women, expressed discomfort, lack of trust, and even fear regarding foreigners. Many mentioned that they were unaware or did not understand the reason for the presence of foreigners or thought that the foreigners were there to carry out deleterious activities such as the sale of blood. For example, participants from all communities told stories about the Amazonian myth of the “pishtacos,” outsiders who killed local residents to sell their blood and fat (22).

When asked specifically about the presence of foreigners providing healthcare support, participants from all communities expressed positive opinions. They specifically mentioned the high quality of the medical equipment and medications provided by foreigners. In addition, participants had advice and requests in order to achieve successful, productive interactions between community members and foreign visitors. They requested that the visitors: 1) provide clear and detailed information about what they plan to do, why, and for whom, 2) ensure that participation is voluntary, and 3) coordinate with community members and authorities. Participants also stated that foreign visitors should speak Spanish, in order to communicate effectively with community members. Finally, one leader underscored the importance of making interventions comprehensive and sustainable over time.

IV. DISCUSSION

We assessed the health of children in four remote rural Amazonian communities as part of a baseline evaluation prior to a short-term humanitarian medical visit. We found a high prevalence of malnutrition, anemia, and intestinal parasites as well as a low rate of complete immunizations compared to rates reported for all the country (4). Residents of all four communities had a similar spectrum of health problems despite differences in the size of the communities, health infrastructure, and access to health services. This was consistent with overall assessments conducted in rural communities of the Amazon Basin (23) and confirms that the sites selected for the HA activity were in need of these activities (24).

The prevalence of stunting in our population was similar to that found in other studies (24, 25), especially those in the Amazon Basin (26-29). We found a negative association between stunting and age, also similar to other studies in Peru and other Latin American countries (26, 28, 30). Our study reported a lower prevalence of anemia compared to others studies in the same age group (24, 28, 31). Among parasites, DNA-based methods demonstrated the absence of *E. histolytica* in this study population. The prevalence of geohelminths and *Giardia lamblia* was similar to those in other studies conducted in Peru, Latin America, Turkey, and Yemen (26, 29, 32-35), in spite of the limitations of the direct observation and a unique stool sample for diagnosis of intestinal parasites. The findings regarding increased prevalence of intestinal polyparasitism and its relation with age were similar to other studies in the Peruvian Highlands with children over five years of age (36). However, a study in Cuba reported that rates of intestinal parasitism were not associated with a child's age (37). All intestinal parasites increased after 1 year of age in our study; a similar correlation with age was found in studies in rural Ecuador and Ethiopia (38, 39).

According to their caretakers, over two-thirds of children between 1 and 4 years had received the full Peruvian immunization schedule. Other studies in resource-limited settings observed similar vaccination rates (40, 41). However, accurate estimation of vaccine coverage can be affected by recall bias and the inability of some caregivers to present their child's immunization card. Many of these vaccines require a cold chain or are given as a multi-dose series, two logistical hurdles that make achieving effective protection in remote communities challenging. Therefore, HA can be a valuable way to deliver single-dose vaccines that require cold storage such as yellow fever, particularly if using non-conventional cold-chain infrastructure. Additionally, HA can serve to extend the coverage of multi-dose vaccines, such as hepatitis B, by including "immunization day" types of activities.

Anemia, intestinal parasites, and impoverished health infrastructure have both immediate and long-term consequences for children's health. The observation of a high prevalence of various health problems in children 1-4 years highlights the importance of targeting health interventions at this group in order to combat their high burden of disease and malnutrition particularly (stunting and anemia). Also including children under 1 year as a target in these interventions could be beneficial for preventing conditions such as stunting, which may be more reversible at that age (42). The non-detection of pediatric malaria is not unusual in this area, where incidence has been relatively low in the recent years (43).

Our mixed-methods study characterized the baseline health needs of the communities and gauged how the local population, including health providers and heads of civil organizations, felt about their current healthcare as well as their opinions of foreign visitors and healthcare personnel. Community members in Lagunas, a launching point for the Pacaya Samiria National Reserve, were not wary of outsiders, whether medical or non-medical, while residents of smaller

communities were concerned about the presence of outsiders in their communities. Nevertheless, participants did not feel threatened by foreign healthcare personnel; instead, they associated these visitors with “better medicine,” “better care,” and “higher technology.” A study conducted in the Democratic Republic of the Congo found similar positive perceptions in the population regarding healthcare providers from humanitarian organizations (44). However, such receptivity is not always present. A study in Sri Lanka showed that some beneficiaries were not completely satisfied with the distribution of goods by NGOs because they felt that HA personnel were not completely transparent and they did not fulfill their expectations, such as providing clean water and sanitation (45). Therefore, HA should always include an assessment of the perspectives of potential direct beneficiaries such as community members and indirect beneficiaries such as healthcare providers (MoH personnel), local authorities, and civil society organizations as a standard procedure for improving the delivery of such activities. Finally, US Air Force personnel and in general personnel delivering HA interventions can also be indirect beneficiaries of the interventions because they improve their clinical skills, as reported in other HA activities (10).

Short-term humanitarian medical visits, like those offered by the US DoD and other organizations, represent an opportunity for populations in remote areas to receive health interventions in a short period of time (46-48). Nevertheless, the challenge is providing care that is needed, effective, and safe. Short-term surgical HAs, focusing on correcting common childhood afflictions such as cleft palate or congenital urologic abnormalities, provide an obvious immediate benefit. However, they have been criticized for a lack of proper supervision of junior medical staff and absence of post-operative follow-up (12). Non-surgical medical HAs present different challenges, such as indiscriminant distribution of pharmaceuticals to a readily accepting population, which may introduce negative effects like antimicrobial resistance, adverse

drug reactions, or iatrogenesis due to a lack of adequate patient background knowledge (14). Also, liberal distribution of medications by foreign entities can also reduce the local population's trust in local healthcare providers. Therefore, comprehensive baseline evaluations, such as the one presented here, are critical for adequately assessing the needs of the local population, obtaining approval for planned activities, and coordinating the participation and support of local healthcare providers. Careful planning and interaction with local, host country partners can help to maximize the benefits of HA and reduce unexpected, negative outcomes (49). Overseas DoD laboratories such as NAMRU-6 are strategically located and can support MEDRETEs with accurate situational awareness, diverse assessment methods and training of local staff. Their capacity building contributions can be valuable to this short-term HA activities and represent non-measured additional benefits of MEDRETEs.

The quantitative data identified several important public health problems in the local communities such as anemia, parasitism, decayed teeth, and low visual acuity. Some of these issues could be solved with one-time HA interventions improved HAs. In addition, we identified indicators such as the prevalence of anemia, prevalence of intestinal parasites, and immunization rates that would provide an easy way to measure the health of the target population before HA is delivered. Moreover, these indicators could be used as a baseline to measure the impact of future HA activities and could serve to establish goals according to characteristics of each HA intervention. The lack of non-intervened communities as a comparison group and the absence of some indicators in our survey such as chronic headaches related to decayed teeth may limit the accurate assessment of the impact of HA activities. However, even with these limitations, a similar post-HA assessment could be extremely valuable in understanding if HA can be a valuable public health tool.

In summary, our baseline visits uncovered two important facts in these four remote communities in the Amazon River basin. First, we noted health deficiencies that may be amenable to short-term healthcare interventions. Second, we perceived openness to medical care provided by personnel from outside the community, including staff from foreign countries. These two observations indicate a possible role for future short-term humanitarian medical visits as an alternative to address the healthcare needs of people living in remote locations, especially vulnerable populations such as children. Our findings highlight the importance of addressing the impact of HA on the inhabitants' perceptions of the local and visiting healthcare personnel. Additional studies need to be conducted to determine whether such visits would result in tangible, objective improvements in health. We intend on carrying out future work to address these questions.

V. REFERENCES

1. Huerga H, Lopez-Velez R. [Infectious diseases in the migrant child]. *Rev Pediatr Aten Primaria*. 2002;4:73-80. Enfermedades infecciosas en el niño inmigrante.
2. World Health Organization. *Removing obstacles to healthy development*. 1999.
3. Hotez PJ, Molyneux DH, Fenwick A, Kumaresan J, Sachs SE, Sachs JD, et al. Control of neglected tropical diseases. *N Engl J Med*. 2007;357(10):1018-27. Epub 2007/09/07.
4. Instituto Nacional de Estadística e Informática (INEI). *DHS Survey Peru: 2000, 2005, 2006, 2007, 2008, 2009*. Lima: Instituto Nacional de Estadística e Informática; 2010.
5. *Analysis of Health Situation of Loreto Region*. Iquitos: Dirección de Epidemiología, Dirección Regional de Salud Loreto; 2010.
6. Hotez PJ, Bottazzi ME, Franco-Paredes C, Ault SK, Periago MR. The neglected tropical diseases of Latin America and the Caribbean: a review of disease burden and distribution and a roadmap for control and elimination. *PLoS Negl Trop Dis*. 2008;2(9):1-11. Epub 2008/09/30.
7. *Sociodemographic profile of the Department of Loreto*. Lima: Instituto Nacional de Estadística e Informática (INEI); 2009.
8. *Peru: Health and nutritional status of mothers and children under five years*. Lima: Instituto Nacional de Estadística e Informática (INEI); 2010.
9. DeCamp M. Scrutinizing global short-term medical outreach. *The Hastings Center report*. 2007;37(6):21-3. Epub 2008/01/09.
10. Birch M, Miller S. Humanitarian assistance: standards, skills, training, and experience. *BMJ*. 2005;330(7501):1199-201. Epub 2005/05/21.
11. Roberts L, Hofmann CA. Assessing the impact of humanitarian assistance in the health sector. *Emerging themes in epidemiology*. 2004;1(1):3. Epub 2005/02/01.

12. Reaves EJ, Schor KW, Burkle FM, Jr. Implementation of evidence-based humanitarian programs in military-led missions: part I. Qualitative gap analysis of current military and international aid programs. *Disaster Med Public Health Prep.* 2008;2(4):230-6. Epub 2008/12/04.
13. Reaves EJ, Schor KW, Burkle FM, Jr. Implementation of evidence-based humanitarian programs in military-led missions: part II. The impact assessment model. *Disaster Med Public Health Prep.* 2008;2(4):237-44. Epub 2008/12/04.
14. Bolton P, Bass J, Murray L, Lee K, Weiss W, McDonnell SM. Expanding the scope of humanitarian program evaluation. *Prehosp Disaster Med.* 2007;22(5):390-5. Epub 2007/12/20.
15. Facchini RE. Humanitarian and civic assistance health care training and cultural awareness promoting health care pluralism. *Mil Med.* 2013;178(5):537-42. Epub 2013/06/13.
16. Oravec GJ, Artino AR, Jr., Hickey PW. Active-duty physicians' perceptions and satisfaction with humanitarian assistance and disaster relief missions: implications for the field. *PLoS One.* 2013;8(3):e57814. Epub 2013/04/05.
17. World Health Organization and UNICEF. Progress on Drinking Water and Sanitation - 2014 update. WHO, Geneva and UNICEF, New York, 2014.
18. Gutierrez Gonzales SC, Arrospide Velasco N. [Manual of laboratory procedures for malaria diagnosis]. In: Ministerio de Salud, Instituto Nacional de Salud, editors. 2003. p. 9-14.
19. Ritchie LS. An ether sedimentation technique for routine stool examinations. *Bull U S Army Med Dep.* 1948;8(4):326. Epub 1948/04/01.
20. Velasquez DE, Arvelo W, Cama VA, Lopez B, Reyes L, Roellig DM, et al. Short report: Molecular insights for *Giardia*, *Cryptosporidium*, and soil-transmitted helminths from a facility-

based surveillance system in Guatemala. *Am J Trop Med Hyg.* 2011;85(6):1141-3. Epub 2011/12/07.

21. Ministerio de Salud del Peru. Resolución Ministerial N°070-2011-MINSA. http://www.minsagobpe/transparencia/dge_normasasp. 2011.

22. de Pribyl R. [Medical anthropology evidences on the Pishtaco origin]. *Rev Peru Med Exp Salud Publica.* 2010;27(1):123-37. Epub 2010/11/13. Evidencias medico antropologicas sobre el origen del Pishtaco.

23. National Monitoring of Indicators for the Follow-up of Reduction of Chronic Malnutrition in Children. Instituto Nacional de Salud (INS); 2004.

24. Status of childhood in Peru. Lima: Instituto Nacional de Estadistica e Informatica (INEI); 2008.

25. Groeneveld IF, Solomons NW, Doak CM. Nutritional status of urban schoolchildren of high and low socioeconomic status in Quetzaltenango, Guatemala. *Rev Panam Salud Publica.* 2007;22(3):169-77. Epub 2007/12/08.

26. Gyorkos TW, Maheu-Giroux M, Casapia M, Joseph SA, Creed-Kanashiro H. Stunting and helminth infection in early preschool-age children in a resource-poor community in the Amazon lowlands of Peru. *Trans R Soc Trop Med Hyg.* 2011;105(4):204-8. Epub 2011/02/26.

27. Muniz PT, Castro TG, Araujo TS, Nunes NB, da Silva-Nunes M, Hoffmann EH, et al. Child health and nutrition in the Western Brazilian Amazon: population-based surveys in two counties in Acre State. *Cad Saude Publica.* 2007;23(6):1283-93. Epub 2007/06/05.

28. Orellana JD, Coimbra CE, Jr., Lourenco AE, Santos RV. Nutritional status and anemia in Surui Indian children, Brazilian Amazon. *J Pediatr (Rio J).* 2006;82(5):383-8. Epub 2006/09/28.

29. Jardim-Botelho A, Brooker S, Geiger SM, Fleming F, Souza Lopes AC, Diemert DJ, et al. Age patterns in undernutrition and helminth infection in a rural area of Brazil: associations with ascariasis and hookworm. *Trop Med Int Health*. 2008;13(4):458-67. Epub 2008/03/04.
30. Victora CG, de Onis M, Hallal PC, Blossner M, Shrimpton R. Worldwide timing of growth faltering: revisiting implications for interventions. *Pediatrics*. 2010;125(3):e473-80. Epub 2010/02/17.
31. Pilger D, Heukelbach J, Diederichs A, Schlosser B, Pereira Leite Costa Araujo C, Keyzers A, et al. Anemia, leukocytosis and eosinophilia in a resource-poor population with helmintho-ectoparasitic coinfection. *J Infect Dev Ctries*. 2011;5(4):260-9. Epub 2011/05/04.
32. De Souza EA, Da Silva-Nunes M, Malafrente Rdos S, Muniz PT, Cardoso MA, Ferreira MU. Prevalence and spatial distribution of intestinal parasitic infections in a rural Amazonian settlement, Acre State, Brazil. *Cad Saude Publica*. 2007;23(2):427-34. Epub 2007/01/16.
33. Kurup R, Hunjan GS. Epidemiology and control of Schistosomiasis and other intestinal parasitic infections among school children in three rural villages of south Saint Lucia. *J Vector Borne Dis*. 2010;47(4):228-34. Epub 2010/12/24.
34. Alyousefi NA, Mahdy MA, Mahmud R, Lim YA. Factors Associated with High Prevalence of Intestinal Protozoan Infections among Patients in Sana'a City, Yemen. *PLoS One*. 2011;6(7):e22044. Epub 2011/07/27.
35. Balcioglu IC, Kurt O, Limoncu ME, Dinc G, Gumus M, Kilimcioglu AA, et al. Rural life, lower socioeconomic status and parasitic infections. *Parasitol Int*. 2007;56(2):129-33. Epub 2007/02/28.
36. Rua O, Romero G, Romani F. Prevalence of intestinal parasitosis in scholar children in a district of the Peruvian highland. *Rev Peru Epidemiol*. 2010;14(2):1-5.

37. Wordemann M, Polman K, Menocal Heredia LT, Diaz RJ, Madurga AM, Nunez Fernandez FA, et al. Prevalence and risk factors of intestinal parasites in Cuban children. *Trop Med Int Health*. 2006;11(12):1813-20. Epub 2006/12/21.
38. Belyhun Y, Medhin G, Amberbir A, Erko B, Hanlon C, Alem A, et al. Prevalence and risk factors for soil-transmitted helminth infection in mothers and their infants in Butajira, Ethiopia: a population based study. *BMC Public Health*. 2010;10:21. Epub 2010/01/21.
39. Jacobsen KH, Ribeiro PS, Quist BK, Rydbeck BV. Prevalence of intestinal parasites in young Quichua children in the highlands of rural Ecuador. *J Health Popul Nutr*. 2007;25(4):399-405. Epub 2008/04/12.
40. Akmatov MK, Mikolajczyk RT. Timeliness of childhood vaccinations in 31 low and middle-income countries. *J Epidemiol Community Health*. 2012;66(7):e14. Epub 2011/05/10.
41. Sinno DD, Shoaib HA, Musharrafieh UM, Hamadeh GN. Prevalence and predictors of immunization in a health insurance plan in a developing country. *Pediatr Int*. 2009;51(4):520-5. Epub 2009/04/30.
42. Grantham-McGregor S, Cheung YB, Cueto S, Glewwe P, Richter L, Strupp B. Developmental potential in the first 5 years for children in developing countries. *Lancet*. 2007;369(9555):60-70. Epub 2007/01/09.
43. Yon C. [Situation of Malaria in the Country]. *Bol Epidemiol (Lima)*. 2010;19(52):1104-6. Situacion de la malaria en el Pais.
44. Dijkzeul D, Wakenge CI. Doing good, but looking bad? Local perceptions of two humanitarian organisations in eastern Democratic Republic of the Congo. *Disasters*. 2010;34(4):1139-70. Epub 2010/07/14.

45. Lee AC. Local perspectives on humanitarian aid in Sri Lanka after the tsunami. *Public Health*. 2008;122(12):1410-7. Epub 2008/08/30.
46. Drifmeyer J, Llewellyn C. Overview of overseas humanitarian, disaster, and civic aid programs. *Mil Med*. 2003;168(12):975-80. Epub 2004/01/15.
47. Drifmeyer J, Llewellyn C. Military training and humanitarian and civic assistance. *Mil Med*. 2004;169(1):23-9. Epub 2004/02/18.
48. Natsios AS. The nine principles of reconstruction and development. *Parameters*. 2005;Autumn:4-20.
49. Maki J, Qualls M, White B, Kleeffeld S, Crone R. Health impact assessment and short-term medical missions: a methods study to evaluate quality of care. *BMC health services research*. 2008;8:121. Epub 2008/06/04.

APPENDICES

Figure 1. Location of study communities in Loreto, Peru.




Table 1. Socioeconomic characteristics of the communities studied

	Saramiriza	P. America	Lagunas	Santa Cruz
Population	1,324	640	5,418	769
Time to Yurimaguas (hours, regular boat)	48	40	16	10
Electricity (hours/day)	3	4	8	4
Improved water sources	Partial	None	Partial	None
Improved sanitation facilities	Partial	None	Partial	None

Table 2. Demographic characteristics of the children studied

	Total (n=457)	Saramiriza (n=85)	P. America (n=25)	Lagunas (n=266)	Santa Cruz (n=81)	<i>p value</i>
Sample*	39%	47%	20%	35%	68%	---
Sex						0.555
Male	49% (224)	45% (38)	40% (10)	50% (133)	53% (43)	
Female	51% (233)	55% (47)	60% (15)	50% (133)	47% (38)	
Age in months**	28.5 ± 18.5	29.6 ± 18.2	34.4 ± 18.8	26.7 ± 18.3	30.9 ± 18.7	0.096
Age groups						0.237
< 1 year	28% (126)	21% (18)	20% (5)	32% (85)	22% (18)	
1 - 2 years	32% (147)	31% (26)	32% (8)	32% (86)	33% (27)	
3 - 4 years	40% (184)	48% (41)	48% (12)	36% (95)	45% (36)	

• This percentage represents the number of children enrolled in total and per community according to the census.

** Mean ± Standard Deviation

Table 3. Nutritional status of children by community and age groups

	Community					<i>p value</i>	Age group			<i>p value</i>
	Total	Saramiriza %(n/N)	P. America %(n/N)	Lagunas %(n/N)	Santa Cruz %(n/N)		< 1 year %(n/N)	1 - 2 years %(n/N)	3 - 4 years %(n/N)	
Anthropometric measures										
Stunting	20% (81/406)	18% (14/77)	33% (7/21)	19% (45/232)	20% (15/76)	0.466	5% (6/110)	25% (33/133)	26% (42/163)	< 0.001
Wasting	3% (13/406)	3% (7/77)	0% (0/21)	3% (8/232)	4% (3/76)	0.929	0% (0/110)	8% (10/133)	2% (3/163)	0.022
Hemoglobin (Hb), g/dL*	11.2 ± 1.1	11.5 ± 1.1	11.2 ± 0.9	11.1 ± 1.1	11.3 ± 1.1	0.028	---	10.8 ± 1.2	11.6 ± 0.9	< 0.001
Anemia % (Hb <11 g/dL)	37% (118/321)	25% (17/67)	32% (6/19)	42% (72/172)	37% (23/63)	0.117	---	55% (77/140)	23% (42/181)	< 0.001

* Mean ± Standard Deviation

Table 4. Prevalence of intestinal parasites of children by community and age groups using microscopy

	Community					<i>p value</i>	Age group			<i>p value</i>
	Total (n=453)	Saramiriza (n=82)	P. America (n=25)	Lagunas (n=265)	Santa Cruz (n=81)		< 1 year (n=126)	1 - 2 years (n=145)	3 - 4 years (n=182)	
Two or more parasites	41%	38%	60%	39%	44%	0.179	6%	32%	71%	< 0.001
One or more parasites	62%	60%	76%	60%	64%	0.439	25%	63%	86%	< 0.001
<i>Ascaris lumbricoides</i>	29%	24%	48%	29%	27%	0.147	6%	28%	46%	< 0.001
<i>Giardia lamblia</i>	20%	30%	32%	16%	17%	0.011	2%	19%	33%	< 0.001
<i>Trichuris trichiura</i>	16%	22%	52%	10%	19%	< 0.001	0%	10%	31%	< 0.001
<i>Entamoeba dispar</i>	8%	9%	0%	5%	19%	0.001	3%	8%	11%	0.034
Hookworms	5%	6%	8%	4%	5%	0.683	1%	2%	9%	< 0.001
<i>Strongyloides stercoralis</i>	2%	2%	12%	1%	4%	0.013	0%	3%	3%	0.081
<i>Cryptosporidium</i> spp.	4%	6%	0%	4%	5%	0.697	2%	8%	4%	0.053
<i>Cyclospora cayetanensis</i>	1%	1%	0%	1%	0%	0.801	1%	1%	1%	1.000

Table 5. Health history of children <5 years

Health issues	% (n/N)
Completed full Peruvian immunization schedule (>1 year)	71 (235/330)
Sickness in the last two weeks:	
- Cough	49 (224/456)
- Fever	34 (155/456)
- Diarrhea	29 (130/456)
When my child had a cough:	
- I took my child to a MoH facility	60 (135/224)
- I sought care or treatment on the first day	37 (82/224)
- My child received care from a physician	45 (101/224)
When my child had a fever:	
- I took my child to a MoH facility	72 (111/155)
- I sought care or treatment on the first day	57 (88/155)
- My child received care from a physician	55 (85/155)
When my child had diarrhea:	
- I took my child to a MoH facility	35 (46/130)
- I sought care or treatment on the first day	26 (34/130)
- My child received care from a physician	25 (32/130)
Treatment for intestinal parasites in the last year	49 (225/456)

MoH = Ministry of Health