



UNIVERSIDAD PERUANA  
**CAYETANO HEREDIA**

EFECTO DE LAS VARIABLES  
ESPACIOTEMPORALES SOBRE LA  
ABUNDANCIA, LA ACTIVIDAD DE  
PICADURA Y LA PARIDAD  
DE *Nyssorhynchus darlingi* (DIPTERA:  
CULICIDAE) EN ZONAS  
PERIURBANAS DE IQUITOS, PERÚ

TESIS PARA OPTAR EL GRADO DE  
MAESTRO EN CONTROL DE  
ENFERMEDADES INFECCIOSAS Y  
TROPICALES

MARLON PIERINO SAAVEDRA ROMERO

LIMA – PERÚ

2025



**ASESOR**

**PhD. GABRIEL CARRASCO ESCOBAR**

**JURADO DE TESIS**

DRA. LARISSA OTERO VEGAS

PRESIDENTE

DR. FELIPE CARLOS YON TORRES

VOCAL

MG. VICTOR ARLES VEGA ZAMBRANO

SECRETARIO (A)

## **DEDICATORIA**

A mis padres, Roberto y Bertha, quienes me formaron como persona, a mis hermanos Alliver, Lady y Darel por apoyarme siempre. A mis amigas y jefas Marta, Manuela, Juliana, Jan y Dionicia por incentivarme a crecer profesionalmente.

## **AGRADECIMIENTO**

A mi familia, por acompañarme incondicionalmente en mi formación y por motivarme siempre a crecer como profesional.

A mis compañeros de campo y buenos amigos del laboratorio ICEMR (International Center of Excellence in Malaria Research) y enfermedades tropicales e infecciosas, en especial a Samantha Solís, Marcos Ríos, Hellen Andrade, Lesly Espinoza, Milagros Ayarza y James Beuzeville, por sus apoyo y profesionalismo, recordando siempre los días de campo.

A mis jefas Marta Moreno, Manuela Herrera, Juliana Morales, Jan Conn y Dionicia Gamboa por su liderazgo en el proyecto, de quienes aprendí mucho como entomólogo, además de su constante motivación a seguir mejorando.

A mis compañeros y catedráticos de la Maestría en control de enfermedades infecciosas y tropicales (MCEIT), en especial a la Doctora Larissa Otero, por sus consejos para la elaboración del protocolo y proyecto de tesis.

A mi asesor Dr. Gabriel Carrasco por su asesoramiento y apoyo en el desarrollo de la presente tesis.

## **FUENTES DE FINANCIAMIENTO.**

Tesis Autofinanciada



## DECLARACIÓN DE ORIGINALIDAD

### Los egresados:

N°	APELLIDOS Y NOMBRES
1.	SAAVEDRA ROMERO MARLON PIERINO

*(Agregar filas adicionales si hay más autores)*

Pertenecientes al programa de la **MAESTRÍA EN CONTROL DE ENFERMEDADES INFECCIOSAS Y TROPICALES**, autores del trabajo titulado: **EFFECTO DE LAS VARIABLES ESPACIOTEMORALES SOBRE LA ABUNDANCIA, LA ACTIVIDAD DE PICADURA Y LA PARIDAD DE Nyssorhynchus darlingi (DIPTERA: CULICIDAE) EN ZONAS PERIURBANAS DE IQUITOS, PERÚ**, el cual ha sido elaborado, sustentado y aprobado, según corresponda, para optar por el grado de **MAESTRO EN CONTROL DE ENFERMEDADES INFECCIOSAS Y TROPICALES** bajo la modalidad de **DESARROLLO TEMÁTICO DE ARTÍCULO**.

En calidad de docentes asesores de la Universidad Peruana Cayetano Heredia:

N°	APELLIDOS Y NOMBRES DEL DOCENTE	FACULTAD	NIVEL DE ASESORÍA
1.	CARRASCO ESCOBAR GABRIEL	FAMED	MAESTRÍA

Declaramos que el contenido del presente documento es original y que las citas y referencias a otros autores cumplen con las normas académicas establecidas. En ese sentido, hacemos constar que:

- El documento presenta un porcentaje de similitud de **9%**, según el reporte emitido por el software **Turnitin®** (identificador de entrega: **2814717578**; fecha de entrega: **14-11-2025**).
- Tras una revisión detallada del reporte y del contenido del trabajo en cuestión, no se han identificado indicios de plagio.
- Se certifica que el documento respeta los principios de integridad académica y cumple con los requisitos institucionales de originalidad.

Lugar y fecha: **Lima, 14 de noviembre de 2025**

Firma del asesor  
N° DNI: 47223865  
ORCID: 0000-0002-6945-0419

Firma del Co-asesor

N° DNI: .....  
ORCID: .....

## ÍNDICE

RESUMEN

ABSTRACT

I.	ARTÍCULO PUBLICADO .....	1
II.	DISCUSIÓN .....	14
III.	REFERENCIAS BIBLIOGRÁFICAS .....	21
IV.	ANEXO (Cuando se trata de un artículo con autor secundario)	

## RESUMEN

**Introducción:** Comprender los factores que influyen en el comportamiento de *Ny. darlingi* es clave para optimizar las intervenciones; este estudio analiza esas variables para aportar evidencia útil a los programas de control. **Metodología:** Entre 2016 y 2017 se realizaron colectas mosquitos con cebo humano en el intra y peri-domicilio en cuatro comunidades amazónicas para evaluar abundancia, picadura horaria, paridad y estacionalidad. **Resultados:** Se capturaron 4,299 *Ny. darlingi*, de los cuales el 80% (3,420) fueron en la temporada de lluvia. La abundancia fue mayor en comunidades ribereñas, Lupuna registró un HBR máximo de 43.8 picaduras-persona en el peri-domicilio. Sta. Emilia fue la única con transmisión activa, alcanzando un EIR de 0.427 en lluvias. En contraste, en comunidades de carretera, aunque la densidad fue baja, se observó una longevidad máxima de 8.7 días y paridad de 100% en marzo en El Triunfo. La mayoría de picaduras ocurrió en el peri-domicilio, principalmente entre las 18:00-21:00 h. En la temporada de lluvias la paridad alcanzó 65%, frente a 37% en seca. **Discusión:** Estos hallazgos evidencian que la transmisión de malaria se concentra en comunidades ribereñas y durante la temporada de lluvias, lo que resalta la necesidad de priorizar intervenciones vectoriales en esos contextos. La actividad exofílica temprana disminuye la eficacia de LLINs e IRS, lo que refuerza la importancia de estrategias complementarias en el peri-domicilio. **Conclusión:** *Ny. darlingi* presentó mayor abundancia y transmisión en comunidades ribereñas, con actividad exofílica temprana y estacionalidad que favorece poblaciones longevas en época de lluvias

**Palabras claves:** *Ny. darlingi*, exofilia, estacionalidad, paridad y control vectorial.

## ABSTRACT

**Introduction:** Understanding the factors that shape the behavior of *Ny. darlingi* is key to optimizing interventions. This study analyzes those variables to provide useful evidence for control programs. **Methods:** From 2016 to 2017, human landing catches were conducted indoors and in peri-domestic areas across four Amazonian communities to evaluate abundance, biting activity, parity, and seasonality. **Results:** A total of 4,299 *Ny. darlingi* were collected, of which 80% (3,420) were during the rainy season. Abundance was higher in riverine communities, with Lupuna recording a maximum HBR of 43.8 bites/person/night in peri-domestic sites. Santa Emilia was the only community with active transmission, reaching an EIR of 0.427 during the rainy season. In contrast, in road-linked communities, although density was lower, El Triunfo showed a maximum longevity of 8.7 days and 100% parity in March. Most bites occurred in peri-domestic areas, especially between 18:00–21:00 h. During the rainy season, parity reached 65% compared to 37% in the dry season. **Discussion:** These findings indicate that *Ny. darlingi* transmission is concentrated in riverine communities and during the rainy season, highlighting the need to prioritize vector interventions in these settings. Early evening exophagic activity reduces the effectiveness of LLINs and IRS, underscoring the importance of complementary peri-domestic strategies. **Conclusions:** *Ny. darlingi* showed greater abundance and transmission in riverine communities, with early exophilic activity marked seasonality favoring long-lived populations during the rainy season.

**Keywords:** *Ny. darlingi*, exophily, seasonality, parity, vector control.

## I. ARTÍCULO PUBLICADO

Bickersmith et al. *Malaria Journal* (2024) 23:112  
<https://doi.org/10.1186/s12936-024-04940-z>

Malaria Journal

### RESEARCH

### Open Access



# Effect of spatiotemporal variables on abundance, biting activity and parity of *Nyssorhynchus darlingi* (Diptera: Culicidae) in peri-Iquitos, Peru

Sara A. Bickersmith<sup>1</sup>, Marlon P. Saavedra<sup>2</sup>, Catharine Prussing<sup>3</sup>, Rachel E. Lange<sup>3</sup>, Juliana A. Morales<sup>2</sup>, Freddy Alava<sup>4</sup>, Joseph M. Vinetz<sup>2,5,6</sup>, Dionicia Gamboa<sup>2,6,7</sup>, Marta Moreno<sup>8\*</sup> and Jan E. Conn<sup>1,3\*</sup>

#### Abstract

**Background** In malaria endemic regions of the Peruvian Amazon, rainfall together with river level and breeding site availability drive fluctuating vector mosquito abundance and human malaria cases, leading to temporal heterogeneity. The main variables influencing spatial transmission include location of communities, mosquito behaviour, land use/land cover, and human ecology/behaviour. The main objective was to evaluate seasonal and microgeographic biting behaviour of the malaria vector *Nyssorhynchus* (or *Anopheles*) *darlingi* in Amazonian Peru and to investigate effects of seasonality on malaria transmission.

**Methods** We captured mosquitoes from 18:00 to 06:00 h using Human Landing Catch in two riverine (Lupuna, Santa Emilia) and two highway (El Triunfo, Nuevo Horizonte) communities indoors and outdoors from 8 houses per community, during the dry and rainy seasons from February 2016 to January 2017. We then estimated parity rate, daily survival and age of a portion of each collection of *Ny. darlingi*. All collected specimens of *Ny. darlingi* were tested for the presence of *Plasmodium vivax* or *Plasmodium falciparum* sporozoites using real-time PCR targeting the small subunit of the 18S rRNA.

**Results** Abundance of *Ny. darlingi* varied across village, season, and biting behaviour (indoor vs outdoor), and was highly significant between rainy and dry seasons ( $p < 0.0001$ ). Biting patterns differed, although not significantly, and persisted regardless of season, with peaks in highway communities at ~20:00 h in contrast to biting throughout the night (i.e., 18:00–06:00) in riverine communities. Of 3721 *Ny. darlingi* tested for *Plasmodium*, 23 (0.62%) were infected. We detected *Plasmodium*-infected *Ny. darlingi* in both community types and most (20/23) were captured outdoors during the rainy season; 17/23 before midnight. Seventeen *Ny. darlingi* were infected with *P. vivax*, and 6 with *P. falciparum*. No infected *Ny. darlingi* were captured during the dry season. Significantly higher rates of parity were detected in *Ny. darlingi* during the rainy season (average 64.69%) versus the dry season (average 36.91%) and by community, Lupuna, a riverine village, had the highest proportion of parous to nulliparous females during the rainy season.

\*Correspondence:

Marta Moreno  
Marta.Moreno@hltm.ac.uk  
Jan E. Conn  
jan.conn@health.ny.gov

Full list of author information is available at the end of the article



©The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Conclusions** These data add a seasonal dimension to malaria transmission in peri-Iquitos, providing more evidence that, at least locally, the greatest risk of malaria transmission is outdoors during the rainy season mainly before midnight, irrespective of whether the community was located adjacent to the highway or along the river.

**Keywords** *Nyssorhynchus darlingi*, Amazonian Peru, Malaria transmission, Vectors, Abundance, Seasonality

## Background

Malaria remains a pressing global health issue despite numerous intervention and treatment efforts, i.e., following a decade-long decrease, the World Health Organization (WHO) reported a worldwide increase of 8 million cases between 2015 and 2017, with an annual death toll estimated at 450,000 persons [1–3]. Overall, in the Americas, Peru contributes annually about 15% of cases [3], and within Peru, Loreto Department in northeastern Amazonian Peru accounts for most (90%) of the country-wide cases [4].

*Nyssorhynchus darlingi* (also known as *Anopheles darlingi*) ([5]) is the main vector of *Plasmodium* transmission in and around the capital of Loreto, Iquitos, since its initial detection there [6] at the beginning of a major malaria epidemic [7]. This mosquito species subsequently spread extensively, likely along river systems [8, 9] through many parts of Amazonian Peru. Transmission hotspots occur mainly in the extensive network of riverine villages and along the Iquitos-Nauta highway, linked frequently to occupation (logging, charcoal production, agriculture, fish farming) [10–12]. Understanding malaria transmission in such diverse community types could help elimination efforts and reduction of the overall malaria case burden in Peru.

*Nyssorhynchus darlingi* is a dominant malaria vector because of its anthropophily, behavioural plasticity [13, 14] and preference for secondary forest and forest fragment edges [15, 16]. Frequently collected biting outdoors in many Peruvian communities [17–19] in some villages in Colombia and Brazil it bites frequently indoors [20, 21]. In general malaria vector biting location (indoors, outdoors) and time, can be modified by the use of IRS (insecticide residual spray) and/or LLINs (long-lasting insecticide nets) [14, 22, 23]. Access to a human blood-meal [18], and environmental factors such as local temperature and humidity, also influence vector biting behaviour. The average infectivity rate of *Ny. darlingi* in Peru ranges from 0.1 to 4% [17, 18, 24], but throughout the Amazon Basin elevated human biting rates (HBR) can contribute to higher-than-expected vectorial capacity [25, 26]. Feeding on humans is common but blood-meal analyses have demonstrated that hosts can include chickens, dogs and cattle, depending on local availability and accessibility [18, 21]. Biting in Peru occurs mainly between dusk and midnight with reported unimodal

and bimodal peaks [9, 14]. The Global Fund Malaria Project PAMAFRO programme [27] was introduced in the Loreto Region, Amazonian Peru, in 2005 for 5 years to reduce malaria transmission. The primary interventions were strengthening of malaria diagnosis and detection, improved malaria case-management, use of insecticide-treated nets (ITNs) and encouragement of community participation in environmental management [28, 29]. The main outcome, by 2010, was the reduction in annual incidence rates from 48.9 to 11.6/1000 [28]. Studies in Loreto prior to the 2005–2010 PAMAFRO initiative detected near-equal proportions of *Ny. darlingi* biting indoors and outdoors [24], but since the end of the PAMAFRO programme, a long-lasting insecticidal net (LLIN) distribution campaign by the Ministry of Health, and the Malaria Cero Programme (MCP) initiated in 2017 [30], several communities have shown increased outdoor-biting [17, 19]. However, in localities where insecticide pressure was relaxed, *Ny. darlingi* again began to bite indoors frequently [14]. This shift to increased indoor biting appears to have resulted from behavioural plasticity, and perhaps also aging of LLINs, as opposed to hypothesized genetic differences in *Ny. darlingi* populations in Loreto [9, 14].

The environment is a powerful driver of mosquito-borne disease prevalence [31, 32]. Throughout much of the Amazon, the greatest risk of malaria transmission is during the rainy season [15, 17, 33], or during the wet-dry transition period [34]. Across the Brazilian Amazon, the length of the rainy season, as well as a range of socio-economic factors, contribute the most to malaria risk [35]. This finding is consistent with earlier observations that rainfall may predict vector abundance [34], although a study in French Guiana determined that relationships between *Ny. darlingi* densities, malaria incidence, rainfall and water level were quite variable, depending on local land-cover and availability of suitable breeding habitat [36]. In Amazonian Peru, rainfall leads to an estimated 10 m increase in river levels and contributes to the generation of larval habitats [15, 37]. Seasonal abundance makes studying vector ecology and behaviour in the dry season difficult as numbers of specimens collected are frequently very low; malaria incidence is also reduced [11].

*Nyssorhynchus darlingi* is highly adapted to anthropogenic landscapes. In Brazil, deforestation patches of ~5 km<sup>2</sup> were significantly correlated with malaria

prevalence [16], partly due to this species' preference for forest fringe habitat [38, 39]. Along the Iquitos-Nauta highway in Peru, Vittor et al. [15] found a 278-fold increase in the Human Biting Rate (HBR) and greater numbers of larval habitats in sites with high deforestation. Despite higher forest coverage in riverine communities compared to those located along highways, Lainhart et al. [9] detected a 3.33-fold higher rate of *Plasmodium* transmission in riverine communities, in addition to higher rates of HBR, infection rates (IR), and entomological inoculation rates (EIR). Understanding habitat-specific influences on *Ny. darlingi* behaviour in anthropogenic landscapes can help predict areas of greatest local risk of *Plasmodium* transmission.

This study aimed to understand the environmental effects of community location and seasonality on mosquito abundance, biting behaviour, and entomological indices linked to malaria transmission of the primary malaria vector, *Ny. darlingi*, in Loreto Department south of Iquitos, Peru. Unlike previous temporal studies of *Ny. darlingi* abundance [9, 14, 17] we were able to include dry season data in addition to the more common rainy season findings from four communities.

## Methods

### Study sites

Adult mosquito collections were conducted in four communities in Loreto Department southwest of Iquitos (3.74°S, 73.25°W) that were each visited six times (except for El Triunfo that was visited seven times), in 2016–2017. Depending on locality, 3–4 collections were undertaken during the rainy season and 2–3 during the dry season (Table 1). Two communities studied are along rivers (Lupuna and Santa Emilia) and two along the Iquitos-Nauta highway (Nuevo Horizonte and El Triunfo) (Fig. 1). These sites have been described elsewhere: Lupuna (LUP) by Moreno et al. [17], Santa Emilia (SEM) by Prussing et al. [14], and Nuevo Horizonte (NHO) and El Triunfo (TRI) by Lainhart et al. [9]. As in Lainhart et al. [9], community habitat assignment was determined by proximity of the settlement to the nearest river: riverine localities were < 1 km and highway localities > 2 km from the nearest river.

### Mosquito sampling

We used the Human Landing Catch (HLC) method to collect adult mosquitoes indoors and outdoors (within five meters of the main house door) during four nights per collection event (a total of twenty-eight nights per community). We randomly chose eight houses and during each consecutive night, we captured mosquitoes from two of these houses. Every collection was conducted from 18:00 to 06:00 h, and collectors rotated every three

hours to account for the effects of individual collector variation on attractiveness to mosquitoes. All mosquitoes captured were separated by hour, trap site location (indoor/outdoor), and by community, and subsequently identified using external morphology at our laboratory in Iquitos by trained personnel using standard keys [40–42]. Specimens were maintained on silica gel at 4 °C until DNA extraction.

### Parity, daily survival rate, and life expectancy

To estimate female age composition of the mosquito population, a proportion (9% or more depending on total numbers captured, except for Jan-2016 in NHO where none were assessed for parity) of females collected were dissected to determine parity rates (PR), daily survival, and age estimation per community per season. Mosquito life expectancy (longevity) in days was calculated by Davidson's method [43]:  $Age = \frac{1}{\log e^P}$ , where  $\ell$  is the natural logarithm of the constant 2.71828 and  $P$  is the probability of a mosquito surviving one day (daily survival rate).  $P$  was calculated as:  $P = \frac{PR}{gc}$  [44], where  $PR$  is the ratio between the number of parous mosquitoes to total number of females dissected, and  $gc$  the duration of the gonotrophic cycle (days). As in Moreno et al. [18], the gonotrophic cycle of 2.19 days was used for rainy season collections, and 2.43 for dry season collections [45].

### Molecular detection of sporozoites in *Ny. darlingi*

Genomic DNA was extracted from each specimen of *Ny. darlingi* using Qiagen DNAeasy blood & tissue kits (Qiagen, Hilden, Germany), and DNA quantification conducted with a Qubit 2.0 Fluorometer (ThermoFisher Scientific, Waltham, MA). Detection of *Plasmodium* infection was conducted using real-time PCR targeting the small subunit of the 18S rRNA, with a triplex TaqMan assay (Life Technologies), as previously described [46]. RT-PCR was conducted on pools of DNA of up to five mosquitoes, in equal DNA concentration, for the presence of *P. vivax* and *P. falciparum*. Specimens from positive pools were tested individually to calculate the infection rate (IR).

### Data analysis

The human biting rate (HBR) was evaluated as the average number of *Ny. darlingi* bites per collector per night. This was based on the total number of *Ny. darlingi* collected. As in Lainhart et al. [9], human biting time patterns were graphed, plotting the average proportion of *Ny. darlingi* collected per hour comparing riverine (LUP and SEM) and highway (NHO and TRI) communities; and then tested for significant differences with the nonparametric Kolmogorov–Smirnov (KS) test in GraphPad Prism version 9.2.0 (Graphpad

**Table 1** Monthly abundance, HBR and EIR of *Nyssorhynchus darlingi* captured indoor and outdoor in Lupuna (LUP), Santa Emilia (SEM), El Triunfo (TRI), and Nuevo Horizonte (NHO), 2016–2017

Community	Mo-Yr	Indoor			Outdoor		
		N	HBR( $\pm$ SE)	EIR	N	HBR( $\pm$ SE)	EIR
LUP	Feb-16	164	20.5(7)	0.000	350	43.8(5.4)	0.000
	Apr-16	154	19.2(3.3)	0.000	328	41(9.7)	0.088
	Jul-16	95	11.9(0.5)	0.000	192	24(4.9)	0.000
	Oct-16	34	4.2(1)	0.000	57	7.1(0.9)	0.000
	Dec-16	27	3.8(0.5)	0.000	43	5.4(0.7)	0.000
	Jan-17	172	21.5(7.5)	0.000	273	34.7(3.4)	0.000
	Subtotal	646			1243		
SEM	Feb-16	152	19(1.2)	0.041	317	39.7(5.1)	0.427
	Apr-16	150	18.8(2.2)	0.000	262	32.8(3)	0.161
	May-16	60	7.5(2)	0.000	127	15.7(3)	0.000
	Sep-16	40	5(1.2)	0.000	84	10.5(1.9)	0.000
	Nov-16	38	4.8(1.1)	0.000	69	8.6(1.2)	0.000
	Jan-17	174	21.8(3)	N/A	283	35.4(3.4)	N/A
	Subtotal	614			1136		
TRI	Jan-16	3	0.4(0.2)	0.000	28	3.5(0.7)	0.000
	Apr-16	7	0.9(0.4)	0.000	18	2.3(1.1)	0.000
	Aug-16	8	1(0.3)	0.000	17	2.1(0.4)	0.000
	Oct-16	5	0.6(0.1)	0.000	13	1.6(0.3)	0.000
	Nov-16	3	0.4(0.1)	0.000	12	1.5(0.4)	0.000
	Mar-17	6	1(0.6)	0.000	21	3.5(1.3)	0.000
	Apr-17	3	0.8(0.8)	0.000	11	2.8(2.3)	0.000
	Subtotal	35			120		
NHO	Jan-16	33	4.7(1)	0.000	80	10(1.7)	0.000
	Apr-16	35	4.4(1.4)	0.000	71	8.9(1.1)	0.000
	Jul-16	29	3.6(0.9)	0.000	32	4(0.6)	0.000
	Oct-16	16	2(0.5)	0.000	23	2.9(0.4)	0.000
	Dec-16	18	2.3(0.6)	0.000	24	3(0.2)	0.000
	Apr-17	49	6.7(1.1)	0.085	95	11.9(1.2)	0.909
	Subtotal	180			325		
	TOTAL	1475			2824		


Mo-Yr: month-year of collection; N: number of *Ny. darlingi* captured; HBR: Human Biting Rate; HBR represents the average bites per person per night (b/p/n) calculated from the mean of collection days within each month/12 h per day per month (2 collectors inside and outside per night); EIR: Entomological Inoculation Rate. EIR not available for SEM January 2017. Italicized lines indicate rainy season collections

Software, San Diego, CA). We calculated the Entomological Inoculation Rate (EIR) by multiplying the HBR by the proportion of *Ny. darlingi* that were determined to be *Plasmodium*-positive by RT-PCR. Sporozoite rates were calculated using the number of positive mosquitoes for *Plasmodium* divided by the total number of mosquitoes tested. To calculate the monthly EIR, we combined the numbers of *P. vivax* and *P. falciparum* and then multiplied the HBR by the proportion of infected specimens per month. Our rationale for combining these two *Plasmodium* species is that even though in the 1990s more *Ny. darlingi* were found to be infected by *P. falciparum* compared with *P. vivax* in

Loreto [24], more recent studies have found the opposite [10, 14, 19, 47].

To test the hypothesis that the peak biting time of *Ny. darlingi* in riverine and highway communities differs, we used average proportions of *Ny. darlingi* collected per hour in the four communities and tested for significance with the KS statistical test [9].

Count data of all *Ny. darlingi* collected in 2016–2017 rainy and dry seasons, across all four collection sites, were analysed (Additional file 1) in RStudio version 1.2.5033 (R 4.0.2; R CORE TEAM). To accommodate the overdispersion in our dataset, negative binomial regression models, using forward and backward stepwise



**Fig. 1** Map of communities (LUP: Lupuna, SEM: Santa Emilia, NHO: Nuevo Horizonte and TRI: El Triunfo) within Loreto Department, Peru, where *Ny. darlingi* were collected for this study. Red rectangle indicates the enlarged Loreto region of Peru. Made with Natural Earth data in QGIS v3

selection for variable selection, were conducted with the MASS package [48]. The `glm.nb()` function, was utilized with the following independent variables: season (dry/rainy), site, biting location (indoor/outdoor), 3-h time period (18:00–21:00, 21:00–24:00, 24:00–03:00, 03:00–06:00), and their interactions. A nonparametric Kruskal–Wallis analysis was also conducted on the count data for comparison to the negative binomial regression results.

Count data of the parous and nulliparous female *Ny. darlingi* in the rainy and dry seasons, across the four collection sites, were analysed (Additional file 2) in RStudio version 1.2.5033 (R 4.0.2; R CORE TEAM). We conducted negative binomial regression models as above but using the following independent variables: season (dry/rainy), site, and 6-h time period (18:00–24:00, 24:00–06:00), and their interactions. Some of the variable categories (site, indoor/outdoor, 3-h time period) were collapsed due to small sample sizes, e.g. of nulliparous females in TRI during the rainy season (see Table 2). The parity rates between community (LUP, SEM, TRI, NHO), season (rainy, dry) and community type (highway,

riverine) were compared using the chi-square test with a statistical significance of  $P < 0.05$  in GraphPad Prism version 9.5.1 for Windows, GraphPad Software (San Diego, California, USA).

## Results

### Mosquito capture data and entomological indices

Overall, 4,330 Anophelinae were collected during this study, of which 4,299 were identified morphologically as *Ny. darlingi*. The rainy season accounted for 3,420 individual *Ny. darlingi* specimens (79.55%), and the dry season for 879 (20.45%). Using ITS2-PCR-RFLP as in Matson et al. [49], we identified one *Nyssorhynchus benarrochi* B individual from SEM. Thirty specimens that were provisionally morphologically identified as non-*Ny. darlingi* failed to amplify, could not be identified molecularly, and were excluded from all analyses.

The HBR of *Ny. darlingi* ranged from 0.4 bites per night (b/p/n) indoors in TRI (highway) to 43.8 b/p/n outdoors in LUP (riverine) (Table 1) and was generally higher during the rainy season (Table 1). The final collection of *Ny.*

**Table 2** Total number, percent and number of nulliparous, parous, and gravid female *Nyssorhynchus darlingi* collected from Lupuna (LUP), Santa Emilia (SEM), El Triunfo (TRI), and Nuevo Horizonte (NHO), 2016–2017, in addition to parity rate, daily survival rate, and life expectancy

Community	Mo-Yr	Total Captured	% Total (N)	% Nulliparous (N)	% Parous (N)	% Gravid (N)	PR	Daily survival rate (P)	Age (days)
LUP	Feb-16	514	32.5 (167)	2.4 (4)	71.9 (120)	25.7 (43)	0.72	0.86	6.63
	Apr-16	482	41.1 (196)	3.5 (7)	28.3 (56)	68.2 (135)	0.28	0.56	1.73
	Jul-16	287	46.3 (133)	4.5 (6)	13.5 (18)	82 (109)	0.14	0.44	1.22
	Oct-16	91	97.8 (89)	5.6 (5)	23.6 (21)	70.8 (63)	0.24	0.55	1.68
	Dec-16	70	98.6 (69)	7.2 (5)	23.2 (16)	69.6 (48)	0.23	0.55	1.66
	Jan-17	445	38.4 (171)	6.4 (11)	60.2 (105)	33.3 (57)	0.60	0.79	4.32
	Subtotal	1889	43.8 (827)	4.6 (38)	40.4 (334)	55 (455)			
SEM	Feb-16	469	9.4 (44)	4.5 (2)	70.5 (31)	25 (11)	0.70	0.85	6.25
	Apr-16	412	10.4 (43)	11.6 (5)	51.2 (22)	37.2 (16)	0.51	0.74	3.27
	May-16	181	16 (29)	17.2 (5)	55.2 (16)	27.6 (8)	0.55	0.76	3.68
	Sep-16	124	16.9 (21)	19 (4)	61.9 (13)	19 (4)	0.62	0.82	5.07
	Nov-16	107	16.8 (18)	16.7 (3)	61.1 (11)	22.2 (4)	0.61	0.82	4.93
	Jan-17	457	10.3 (47)	12.8 (6)	46.8 (22)	40.4 (19)	0.47	0.71	2.88
	Subtotal	1750	11.5 (202)	12.4 (25)	56.9 (115)	30.7 (62)			
TRI	Jan-16	31	54.8 (17)	0 (0)	100 (17)	0 (0)	1.00	1.00	N/A
	Apr-16	25	80 (20)	5 (1)	60 (12)	35 (7)	0.60	0.79	4.29
	Aug-16	25	100 (25)	20 (5)	36 (9)	44 (11)	0.36	0.66	2.38
	Oct-16	18	100 (18)	22.2 (4)	38.9 (7)	38.9 (7)	0.39	0.68	2.57
	Nov-16	15	100 (15)	20 (3)	40 (6)	40 (6)	0.40	0.69	2.65
	Mar-17	27	33.3 (9)	0 (0)	77.8 (7)	22.2 (2)	0.78	0.89	8.71
	Apr-17	14	50 (7)	0 (0)	100 (7)	0 (0)	1.00	1.00	N/A
Subtotal	155	71.6 (111)	11.7 (13)	58.6 (85)	29.7 (33)				
NHO	Jan-16	113	0 (0)	N/A (0)	N/A (0)	N/A (0)	N/A	N/A	N/A
	Apr-16	106	53.8 (57)	12.3 (7)	59.6 (34)	28.1 (16)	0.60	0.79	4.24
	Jul-16	61	100 (61)	26.2 (16)	37.7 (23)	36.1 (22)	0.38	0.67	2.49
	Oct-16	39	100 (39)	35.9 (14)	30.8 (12)	33.3 (13)	0.31	0.62	2.06
	Dec-16	42	100 (42)	23.8 (10)	38.1 (16)	38.1 (16)	0.38	0.67	2.52
	Apr-17	144	44.4 (64)	12.5 (8)	59.4 (38)	28.1 (18)	0.59	0.79	4.20
	Subtotal	505	52.1 (263)	20.9 (55)	46.8 (123)	32.3 (85)			
TOTAL	4299	32.6 (1403)	9.3 (131)	45.4 (637)	45.3 (635)				


<sup>a</sup> % Total percent of females dissected for parous status of total captured

*darlingi* in January 2017 from SEM was not analysed for *Plasmodium* for logistical reasons, thus a total of 3,721 *Ny. darlingi* were tested for *Plasmodium*. Of these, 23 (0.62%) were infected. In the localities where *Plasmodium*-infected *Ny. darlingi* were detected (Fig. 2), the EIR ranged from a high of 0.909 outdoors in NHO (highway) to a low of 0.041 indoors in SEM (riverine) (Table 1). No infected *Ny. darlingi* were collected during the dry season.

#### Parity, daily survival rate, and life expectancy

We dissected 1,403/4,299 or 33% of the total number of *Ny. darlingi* from the four localities (Table 2). There were

637 parous, 635 gravid and 131 nulliparous *Ny. darlingi* detected (Table 2 and Additional file 2: Fig. S1). The PR in the rainy season ranged from 0.28 in LUP—1.00 in TRI whereas during the dry season it was 0.14 in LUP—0.62 in SEM. In the four localities, the average estimated mosquito age was somewhat higher during the rainy season (range 1.73–8.71 days) compared with the dry season (range 1.22–5.07 days), and the highest mosquito age was detected in TRI during the rainy season (8.71 days). There was a significant difference in the proportion of parous mosquitoes by site ( $\chi^2=8.04$ ,  $df=3$ ,  $P=0.0452$ ) as well as season ( $\chi^2=8.621$ ,  $df=1$ ,  $P=0.0033$ ), but not for community type ( $\chi^2=3.724$ ,  $df=1$ ,  $P=0.0536$ ).



**Fig. 2** Abundance of *Ny. darlingi* collected indoors vs outdoors and rainy vs dry season in four collection sites in Loreto, Peru, 2016–2017. Collection times of infected specimens are indicated by a closed circle (*P. vivax*) or a closed triangle (*P. falciparum*)

#### Seasonal, location (indoor/outdoor) and community effects

In general, more *Ny. darlingi* were captured during the rainy season compared with the dry season. Regardless of community or season, more *Ny. darlingi* were captured outdoors than indoors from all four sites; more were captured in riverine (LUP, SEM) than highway (TRI, NHO) communities (Table 1; Fig. 2). Regardless of location, season or community type, *Ny. darlingi* bit throughout the night, although there was a more pronounced early evening peak in the highway communities of TRI and NHO between 19:00 and 21:00 compared with LUP and SEM (Additional file 2: Figs. S2 and S3). There were no significant differences detected in biting patterns by the KS test ( $p=0.8475$ ; Fig. 3).

Results from negative binomial regression indicated significant differences in counts across sites, season, time period, and between indoor and outdoor captures (Table 3). Compared to the three other collection sites, significantly more *Ny. darlingi* were captured in LUP. In addition, more mosquitoes were collected during the rainy than dry season, outdoors versus indoors, and during the first time period (18:00–21:00) across all sites. Significant interactions (time period X indoor/outdoor, season X time period, season X site, and site X indoor/outdoor) suggest that these relationships are contextually dependent. A few of these interactions were significant only for TRI in comparison to LUP: season X site (indicating a lower rainy: dry season ratio in TRI than


the other sites) and site X indoor/outdoor (indicating a higher outdoor: indoor ratio in TRI than the other sites). A Kruskal–Wallis analysis of the same *Ny. darlingi* count data supported the negative binomial regression findings (Supplemental Table 1).

#### Malaria incidence data

During our study (2016–2017), the Annual Parasite Index (API), based on microscopic identification of blood smears, ranged from a low of 37.8 (TRI 2016) to a high of 504.1 in LUP (Table 4). In 2017, two of the communities registered a lower API compared with 2016: TRI, population 238, increased marginally from 37.8 to 54.6 and SEM, population 204, increased from 181.4 to 279.4. In the four communities during the time-frame of the present study (2016–2017) more malaria cases (~65%) were *P. vivax* (~65%) compared with *P. falciparum* (35%). These values are similar to national Peruvian averages for 2016–2017 [28, 50].

#### Discussion

Results are congruent with previous studies in Amazonian Peruvian of *Ny. darlingi* that found the highest risk of acquiring malaria to be outdoors before midnight during the rainy season. Our new data underscore the low risk of local malaria transmission during the dry season, due in part to a highly significant reduction in *Ny. darlingi* abundance together with significantly lower parity rates, compared with the rainy season. In agreement



**Fig. 3** Average proportion of *Ny. darlingi* collected hourly by community type (LUP and SEM, riverine; NHO and TRI, highway). Error bars represent 95% confidence intervals

with previous studies in the peri-Iquitos region of Loreto Department in Amazonian Peru [10, 19], *Ny. darlingi* was the predominant anopheline species collected in the present study (99.3%). The HBR, IR and EIR were all within the range of other similar regional studies that have focused mainly on *Ny. darlingi* (Table 1).

In peri-Iquitos, *Ny. benarrochi* B appears to be relatively uncommon, as previous reports demonstrate [14, 19, 51, 52]. In some regions, notably Datem del Marañon province, it was [8] and remains [53, 54] highly abundant. It is a regional or secondary *Plasmodium* vector in southern Colombia, eastern Peru [55], eastern (Amazonian) Ecuador [53] and Datem del Marañon [54].

Despite high net distribution and cover and access (many residents in the present study use both *tocuyos* and LLINs as we have shown previously [17, 19], *Plasmodium* continues to be transmitted and the APLs, especially in the two riverine villages of LUP and SEM, were high in both 2016 and 2017 (Table 4). In similar villages in Mazan district, Loreto region, Peru, we demonstrated that during the early evening (17:00–20:00), between 20 and ~80% of the human population was protected under nets, leaving adequate hosts available for mosquitoes to obtain a blood meal [19]. The most likely explanation for transmission in the present study is that some of the

residents were not using their bednets during the early evening hours and therefore not protected. We observed in this study as well as in [19], that residents indoors were occupied eating or watching television, not using their nets, or they were outdoors bathing in a nearby river or playing soccer. It has been suggested that interventions that are more in synch with the malaria endemic community lifestyles and can be incorporated into routine activities will have a better chance of protecting individuals [56].

Notably, the riverine village of LUP has persisted as a malaria transmission hot spot for several years [28]. In the present study, the highest *Ny. darlingi* HBR was reported in LUP and there were significantly more *Ny. darlingi* collected in this village compared with the other three communities (Tables 1 and 3). The possible explanations for the complex malaria transmission scenario in LUP are varied: (1) the high biting rate may be an adaptation to increase *Plasmodium* transmission as infectivity rates in *Ny. darlingi* are relatively low [25]; (2) our prior larval sampling in LUP revealed that a slow-moving stream traversing the village has been a productive and permanent breeding site for *Ny. darlingi* [37]; (3) high *Plasmodium* genetic diversity and gene flow among geographically relatively far-flung communities including

**Table 3** Negative binomial regression model of *Nyssorhynchus darlingi* counts collected in four sites (Lupuna, Nuevo Horizonte, Santa Emilia, El Triunfo) during rainy and dry seasons

Variable	$\beta$	$e^{\beta}$	SE	p value
Intercept	1.71	5.60	0.11	< 0.0001
Site (ref=Lupuna)				
NHO	-1.13	0.32	0.14	< 0.0001
SEM	-0.27	0.76	0.13	<b>0.0372</b>
TRI	-2.49	0.08	0.22	< 0.0001
Season (ref=dry)	1.17	3.23	0.11	< 0.0001
Time period (ref=18:00–21:00)				
21:00–24:00	-1.34	0.26	0.16	< 0.0001
24:00–03:00	-0.66	0.52	0.14	< 0.0001
03:00–06:00	-0.50	0.61	0.14	< 0.0001
Indoor/outdoor (ref=indoor)	0.56	1.76	0.11	< 0.0001
Time period x indoor/outdoor				
21:00–24:00 outdoor	0.53	1.70	0.14	< 0.0001
24:00–03:00 outdoor	0.00	1.00	0.14	0.9973
03:00–06:00 outdoor	-0.14	0.87	0.14	0.3190
Season x time period				
Rainy x 21:00–24:00	0.34	1.41	0.15	<b>0.0248</b>
Rainy x 24:00–03:00	0.02	1.02	0.14	0.8997
Rainy x 03:00–06:00	-0.42	0.66	0.14	<b>0.0037</b>
Season x site				
Rainy NHO	-0.24	0.79	0.15	0.1049
Rainy SEM	0.05	1.05	0.13	0.7085
Rainy TRI	-0.74	0.48	0.20	<b>0.0002</b>
Site x indoor/outdoor				
NHO outdoor	-0.06	0.94	0.14	0.6710
SEM outdoor	-0.01	0.99	0.12	0.9037
TRI outdoor	0.60	1.83	0.22	<b>0.0062</b>

$\beta$ : regression coefficient;  $e^{\beta}$ : exponentiated regression coefficient (z-value); SE: Standard error

Significance level  $p < 0.05$  in bold

LUP and SEM facilitates the movement of malaria parasites [57]; (4) recurrent seasonal flooding adds to the abundance and fluctuations of *Ny. darlingi* populations; and (5) there is transmission microheterogeneity within the village of LUP [58]. It is difficult to quantify the contribution of each of these factors but integrating multiple

effective interventions such as regular distribution of LLINs and/or impregnated hammocks and hammock nets, (after testing for effectiveness in reduction of indoor and outdoor transmission) to every resident; detection and monthly treatment of larval breeding sites (as in [59]); and routine detection and treatment of both symptomatic and asymptomatic/sub-microscopic *Plasmodium* carriers [60] would undoubtedly reduce malaria transmission. A comparison of human blood samples tested for presence of *Plasmodium* spp. in Santa Emilia by microscopy versus PCR by Ramirez [61] underscore the importance of testing and treatment of both symptomatic and asymptomatic/sub-microscopic *Plasmodium* carriers to cut transmission. A multi-pronged programme (PAMAFRO) was successful in dramatically reducing malaria in Amazonian Peru from 2005 to 2010 [28]. The Plan Maria Cero in Loreto, implemented from 2017 to 2020, reduced malaria cases by 75% [62]. Subsequently, in 2022, the national malaria elimination plan was launched with the main objective of reducing malaria cases in Peru by 90% by 2030 [63].

The propensity of the primary regional vector *Ny. darlingi* to bite outdoors in the early evening, at least in many riverine communities in the Amazon [64, 65] when residents are outdoors eating, relaxing, or working, constitutes a major coverage gap [19]. One plausible vector control intervention for local communities with housing that frequently includes incomplete or fewer than four walls could be the use of eave ribbons impregnated with a mosquito repellent such as those that have been shown to be highly effective in several sub-Saharan malaria endemic countries [66–68]. Clearly, such interventions would need to be tested rigorously in the Latin American context.

We were able to collect adequate sample sizes during the dry season for analysis in this study ( $n = 879$  (20.45%)) and detected significant seasonal differences in abundance (Table 3) and parity. The lack of dry season *Ny. darlingi* infected with *Plasmodium* supports earlier findings that the risk for transmission in this region is during the rainy season, although farther west, in Datem del Marañon province, even in the dry season there is a risk of acquiring malaria [54]. There are two probable reasons

**Table 4** Number of malaria cases in the four sampled communities in Loreto, Peru (2016 and 2017)

Community	Population	No. cases 2016	API 2016	No. cases 2017	API 2017
Lupuna	365	184	504.1	163	446.6
Santa Emilia	204	37	181.4	57	279.4
El Triunfo	238	9	37.8	13	54.6
N. Horizonte	375	61	162.7	44	117.3

N: Nuevo; Data from Ministry of Health, Iquitos, Peru; API: Annual Parasite Index calculated as number of confirmed malaria cases per 1,000 individuals

for this. First, during the low-transmission (dry) season, despite a substantial burden of sub-microscopic infections in Loreto [12, 61] and the fact that persons with submicroscopic malaria can infect anophelines [69], generally very few mosquitoes are infected at low gametocyte densities with standard membrane feeders and transmission is considered much less likely to occur [60, 70]. Nevertheless, a comparison of microscopy versus real time PCR results of *Plasmodium*-positive residents from the community of Santa Emilia from January–September 2016, demonstrated a very biologically significant difference i.e., 6.5% (92/1416) infectivity by microscopy and 24.34% (295/1212) by PCR. The data support the consideration of testing and treating asymptomatic inhabitants, particularly of malaria hotspots.

The second reason is that malaria transmission occurs not only in villages but also in temporary locations (for example, logging or mining camps) linked to seasonal occupation [10, 12]. We did not collect anopheline specimens from such localities to test for this study but the Parker et al. [10] study along the Mazan River during both rainy and dry seasons detected two of 967 *Ny. darlingi* specimens infected, one positive for *P. falciparum* and one for *P. vivax*, from occupation-related logging sites.

The peak biting patterns for *Ny. darlingi* reported here are similar to those in Lainhart et al. [9], in that they differ between riverine and highway communities, albeit not significantly. Thus, in the present study, there was no significant difference between seasonal patterns, and we reject our initial hypothesis of differences between village types (riverine and highway) and instead propose that these biting patterns result from the spatiotemporal availability of human (and possibly other animal and/or bird) hosts [9, 18, 71, 72].

Our parity data indicate that the majority of *Ny. darlingi* females seeking a human bloodmeal are parous, i.e., older, as previous studies have demonstrated in Amazonian Peru [18] and Brazil [45] and, therefore, they could potentially be infected with *Plasmodium*. However, only during the March collection in TRI were samples of *Ny. darlingi* old enough (range 7.24–9.13 days) to sustain the *P. vivax* sporogonic cycle, calculated with the Moshkovsky method in Barros et al. [45]). Parity rates were higher for *Ny. darlingi* during peak transmission (second half of the rainy season and beginning of the dry season) in LUP, NHO and TRI, although this does not hold for SEM (Table 2). Limitations of this study include the high variability in the range of mosquitoes that were dissected and analysed to estimate parity among sites and seasons, as well as in some cases, very low numbers of nulliparous mosquitoes, such that the vector age could not be calculated. Dissections can be inaccurately interpreted

and training in the apparently simple technique is essential. Determination of age has long been a vexing issue in vector biology but a promising new surveillance method, based on deep learning of mid-infrared spectra of mosquito cuticle, was able to accurately and cost-effectively identify both species and age class among three closely related Africa vectors, *Anopheles gambiae*, *Anopheles arabiensis* and *Anopheles coluzzii* [73]. Hopefully, this method can soon be applied to Latin American and other regional malaria vectors and make a significant impact on malaria transmission reduction.

## Conclusions

Malaria incidence in Peru is highest when the interactions between the various ecological and human factors are optimal for malaria transmission [28] and may be region specific. For example, in Roraima state, Brazil, peak malaria incidence and the highest parity rates occurred during the dry season [45], whereas in southern Venezuela, the peak malaria incidence occurred one month after peak biting rates by *Ny. darlingi* and *Nyssorhynchus marajoara*, and there was no correlation with rainfall [74]. In the present study, malaria transmission is optimal during the rainy season when there is an abundance of breeding sites, parous female *Ny. darlingi*, and available human hosts, especially outdoors during the early evening hours that coincide with the peak human biting rates of *Ny. darlingi*.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12936-024-04940-z>.

**Additional file 1: Count dataset.** *Nyssorhynchus darlingi* by collection site, season (rainy/dry), biting location (indoor/outdoor), and time period for all four collection sites (LUP, NHO, SEM, TRI).

**Additional file 2: Table S1.** Kruskal-Wallis analysis on ranked abundance of *Nyssorhynchus darlingi* in four collection sites (Lupuna, Nuevo Horizonte, Santa Emilia, El Triunfo), during rainy and dry seasons 2016–2017.

**Figure S1.** Average parity rate for each collection site comparing: **A** Before vs. after midnight collections; **B** Indoor vs. outdoor collections; **C** Rainy vs. dry collections. **Figure S2.** Average proportion of *Ny. darlingi* collected hourly biting indoor vs. outdoor for each collection site. Confidence intervals not shown for clarity. **Figure S3.** Average proportion of *Ny. darlingi* collected hourly biting by season for each collection site. Confidence intervals not shown for clarity.

## Acknowledgements

We thank all study participants and the leaders and residents of Lupuna, Santa Emilia, El Triunfo, and Nuevo Horizonte for agreeing to host this study in their communities and all field workers (David Arimuya, Abraham Wichez, Mario Flores, Delfin Mamerto, Gloria Rodriguez, Santiago Ruiz, Papa Segundo, Papa Wilson and Victor Aquibuar) for their dedication during the seasonal field collections. We also thank all the people involved in the Amazon ICEMR project in Lima and Iquitos. We are grateful to Dirección Regional de Salud (DIRESA, Iquitos, Loreto) for collaboration and facilitating logistics in Loreto Department. This publication has been possible thanks to the authorization and permits (no. 0424-2012-AG-DGFFS-DGEFFS) from Dirección de Gestión Forestal y de

Fauna Silvestre y la Dirección General Forestal y de Fauna Silvestre del Ministerio de Agricultura de la República del Perú.

#### Author contributions

CP, MM and JEC conceived and designed the study. CP, MM and MPS supervised the study. FA provided guidance on field site selection and malaria case data for each community. CP, MM and MPS conducted the study. SAB analysed the data with help from RL and MPS. SAB, RL and JEC wrote the manuscript. All authors read and approved the final manuscript.

#### Funding

This project was funded by NIH-NIAID (U19AI089681) to JMV and NIH-NIAID (R01AI110112) to JEC. The Biodefense and Emerging Infectious Disease training fellowship Grant T32AI05532901 provided partial support for CP.

#### Availability of data and materials

The data supporting the conclusions of this article are included within the article. The raw data used and/or analysed in this study are available from the corresponding author upon reasonable request.

#### Declarations

##### Ethics approval and consent to participate

Study protocols were approved by the Ethics Review Board of the Regional Health Direction of Loreto (477-2016), Universidad Peruana Cayetano Heredia, in Lima (184-09-16), Asociación Benéfica PRISMA, Lima, Peru and the Human Subjects Protection Program of the University of California San Diego, La Jolla, CA, USA. The New York State Department of Health Institutional Review Board considers mosquito collecting to be a Risk Management issue. Collections were only conducted by trained professionals and standard precautions were taken by all collectors.

##### Consent for publication

Not applicable.

##### Competing interests

The authors declare that they have no competing interests.

##### Author details

<sup>1</sup>Wadsworth Center, New York State Department of Health, Albany, NY, USA. <sup>2</sup>Laboratorio ICENR-Amazonia, Laboratorios de Investigación y Desarrollo, Facultad de Ciencias e Ingeniería, Universidad Peruana Cayetano Heredia, Lima, Peru. <sup>3</sup>Department of Biomedical Sciences, School of Public Health, State University of New York-Albany, Albany, NY, USA. <sup>4</sup>Gerencia Regional de Salud de Loreto (GERESA), Iquitos, Peru. <sup>5</sup>Section of Infectious Diseases, Department of Internal Medicine, Yale School of Medicine, New Haven, CT, USA. <sup>6</sup>Instituto de Medicina Tropical "Alexander Von Humboldt", Universidad Peruana Cayetano Heredia, Lima, Peru. <sup>7</sup>Departamento de Ciencias Celulares y Moleculares, Facultad de Ciencias e Ingeniería, Universidad Peruana Cayetano Heredia, Lima, Peru. <sup>8</sup>Department of Infection Biology, London School of Hygiene & Tropical Medicine, London, UK.

Received: 17 August 2023 Accepted: 10 April 2024  
Published online: 19 April 2024

#### References

- World Health Organization. World Malaria Report 2016. Geneva: World Health Organization; 2016. p. 148.
- World Health Organization. World Malaria Report 2017. Geneva: World Health Organization; 2017. p. 196.
- World Health Organization. World Malaria Report 2018. Geneva: World Health Organization; 2018. p. 210.
- World Health Organization. World malaria report 2020: 20 years of global progress and challenges. Geneva, Switzerland: World Health Organization; 2020. p. 247.
- Foster PG, de Oliveira TMP, Berigo ES, Conn JE, Sant'Ana DC, Nagaki SS, et al. Phylogeny of Anophelinae using mitochondrial protein coding genes. *R Soc Open Sci*. 2017;4:170758.
- Fernández R, Carbajal F, Quintana J, Chauca H, Watts DM, de Presencia A. *INI darlingi* (Diptera: Culicidae), en alrededores de la ciudad de Iquitos, Loreto-Perú. *Boletín de la Soc Peruana de Enfermedades Infecciosas y Trop*. 1996;5:10–20.
- Aramburo-Guarda J, Ramal Assayag C, Witzig R. Malaria reemergence in the Peruvian Amazon region. *Emerg Infect Dis*. 1999;5:209–15.
- Schoeler GB, Flores-Mendoza C, Fernandez R, Davila JR, Zyzak M. Geographical distribution of *Anopheles darlingi* in the Amazon Basin region of Peru. *J Am Mosq Control Assoc*. 2003;19:286–96.
- Lainhart W, Bickersmith S, Nadler K, Moreno M, Saavedra M, Chu VM, et al. Evidence for temporal population replacement and the signature of ecological adaptation in a major Neotropical malaria vector in Amazonian Peru. *Malar J*. 2015;14:375.
- Parker BS, Paredes Dlonogui M, Penataro Yori P, Escobedo K, Florin D, Rengifo Pinedo S, et al. Hyperendemic malaria transmission in areas of occupation-related travel in the Peruvian Amazon. *Malar J*. 2013;12:178.
- Rosas-Aguirre A, Speybroeck N, Llanos-Cuentas A, Rosanas-Urgell A, Carrasco-Escobar G, Rodríguez H, et al. Hotspots of malaria transmission in the Peruvian Amazon: rapid assessment through a parasitological and serological survey. *PLoS ONE*. 2015;10:e0137458.
- Carrasco-Escobar G, Gamboa D, Castro MC, Bongdiwala SI, Rodríguez H, Contreras-Mancilla J, et al. Micro-epidemiology and spatial heterogeneity of *P. vivax* parasitaemia in riverine communities of the Peruvian Amazon: a multilevel analysis. *Sci Rep*. 2017;7:8062.
- Vezzenegho SB, Adde A, Pommier de Santi V, Issaly J, Carinci R, Gabork P, et al. High malaria transmission in a forested malaria focus in French Guiana: how can esophageal *Anopheles darlingi* thwart vector control and prevention measures? *Mem Inst Oswaldo Cruz*. 2016;111:561–9.
- Prussing C, Moreno M, Saavedra MP, Bickersmith SA, Gamboa D, Alava F, et al. Decreasing proportion of *Anopheles darlingi* biting outdoors between long-lasting insecticidal net distributions in peri-Iquitos, Amazonian Peru. *Malar J*. 2018;17:86.
- Vittor AY, Gilman RH, Tielich J, Glass G, Shields T, Lozano WS, et al. The effect of deforestation on the human-biting rate of *Anopheles darlingi*, the primary vector of falciparum malaria in the Peruvian Amazon. *Am J Trop Med Hyg*. 2006;74:3–11.
- Chaves LSM, Conn JE, López RMM, Salim MAM. Abundance of impacted forest patches less than 5 km<sup>2</sup> is a key driver of the incidence of malaria in Amazonian Brazil. *Sci Rep*. 2018;8:7077.
- Moreno M, Saavedra MP, Bickersmith SA, Lainhart W, Tong C, Alava F, et al. Implications for changes in *Anopheles darlingi* biting behaviour in three communities in the peri-Iquitos region of Amazonian Peru. *Malar J*. 2015;14:290.
- Moreno M, Saavedra MP, Bickersmith SA, Prussing C, Michalski A, Tong Rios C, et al. Intensive trapping of blood-fed *Anopheles darlingi* in Amazonian Peru reveals unexpectedly high proportions of avian blood-meals. *PLoS Negl Trop Dis*. 2017;11:e0055337.
- Saavedra MP, Conn JE, Alava F, Carrasco-Escobar G, Prussing C, Bickersmith SA, et al. Higher risk of malaria transmission outdoors than indoors by *Nyssorhynchus darlingi* in riverine communities in the Peruvian Amazon. *Parasit Vectors*. 2019;12:374.
- de Almeida NCV, Louzada J, Neves M, Carvalho TM, Castro-Ahues J, Silveira-Nascimento TF, et al. Larval habitats, species composition and distribution of malaria vectors in regions with autochthonous and imported malaria in Roraima state, Brazil. *Malar J*. 2022;21:13.
- Piedrahíta S, Álvarez N, Naranjo-Díaz N, Bickersmith S, Conn J, Correa M. *Anopheles* blood meal sources and entomological indicators related to *Plasmodium* transmission in malaria endemic areas of Colombia. *Acta Trop*. 2022;233:106567.
- Muslimi AK, Smith DL, Kilama M, Rek J, Arinaitwe E, Nankabinwa J, et al. Impact of vector control interventions on malaria transmission intensity, outdoor vector biting rates and *Anopheles* mosquito species composition in Tororo, Uganda. *Malar J*. 2019;18:445.
- Bugoro H, In'ofa C, Mackenzie DG, Apakramo A, Hevabo W, Corcoran S, et al. Changes in vector species composition and current vector biology and behaviour will favour malaria elimination in Santa Isabel Province, Solomon Islands. *Malar J*. 2011;10:287.

24. Reinbold-Wasson DD, Sardelis MR, Jones JW, Watts DM, Fernandez R, Carbajal F, et al. Determinants of *Anopheles* seasonal distribution patterns across a forest to periurban gradient near Iquitos, Peru. *Am J Trop Med Hyg*. 2012;86:459–63.
25. Sallum MAM, Conn JE, Berço ES, Laporta GZ, Chaves LSM, Bickersmith SA, et al. Vector competence, vectorial capacity of *Nyssorhynchus darlingi* and the basic reproduction number of *Plasmodium vivax* in agricultural settlements in the Amazonian Region of Brazil. *Malar J*. 2019;18:117.
26. Zimmerman RH, Galardo AKR, Lounibos LP, Galardo C, Bahar AK, van Santen E. Vectorial capacities for malaria in eastern Amazonian Brazil depend on village, vector species, season, and parasite species. *Malar J*. 2022;21:237.
27. Rosas-Aguilera A, Guzman-Guzman M, Moreno-Gutierrez D, Rodriguez-Ferrucci H, Vargas-Pacherez D, Acuna-Gonzalez Y. Long-lasting insecticide-treated bednet ownership, retention and usage one year after their distribution in Loreto, Peru. *Rev Peru Med Exp Salud Publica*. 2011;28:228–36.
28. Soto-Calle Y, Rosas-Aguilera A, Llanos-Cuentas A, Abath E, DeDeken R, Rodriguez H, et al. Spatio-temporal analysis of malaria incidence in the Peruvian Amazon Region between 2002 and 2013. *Sci Rep*. 2017;7:40350.
29. Janke MM, Recalde-Coronel GC, Damasceno CP, Salmón-Mulanovich G, Barbieri AF, Lescano AG, et al. The impact of sustained malaria control in the Loreto region of Peru: a retrospective, observational, spatially-varying interrupted time series analysis of the PMAAPRO program. *Lancet Reg Health Am*. 2023;20:100477.
30. Sanchez-Castro EE, Cahuana GM, Garcia-Rios CJ, Guerra-Duarte C, Chauca P, Tapia-Limonchi R, et al. Health and economic burden due to malaria in Peru over 30 years (1990–2019): findings from the global burden of diseases study 2019. *Lancet Reg Health Am*. 2022;15:100347.
31. Norris DE. Mosquito-borne diseases as a consequence of land use change. *EcolHealth*. 2004;1:19–24.
32. Shocket MS, Anderson CB, Caldwell JM, Childs ML, Couper LI, Han S, et al. Environmental drivers of vector-borne diseases. In: Drake JM, Borsari M, Strand M, editors, et al., *Population biology of vector-borne diseases*. Oxford: Oxford University Press; 2020.
33. da Silva-Vasconcelos A, Kato MY, Mourao EN, de Souza RT, Lacerda RN, Sibajev A, et al. Biting indices, host-seeking activity and natural infection rates of anopheline species in Boa Vista, Roraima, Brazil from 1996 to 1998. *Mem Inst Oswaldo Cruz*. 2002;97:151–61.
34. Galardo AK, Zimmerman RH, Lounibos LP, Young LJ, Galardo CD, Aruda M, et al. Seasonal abundance of anopheline mosquitoes and their association with rainfall and malaria along the Matapi River, Amapa, Brazil. *Med Vet Entomol*. 2009;23:335–49.
35. Canelas T, Castillo-Salgado C, Baquero OS, Ribeiro H. Environmental and socioeconomic analysis of malaria transmission in the Brazilian Amazon, 2010–2015. *Rev Saude Publica*. 2019;53:49.
36. Girod R, Roux E, Berger F, Stefani A, Gaborit P, Carinci R, et al. Unravelling the relationships between *Anopheles darlingi* (Diptera: Culicidae) densities, environmental factors and malaria incidence: understanding the variable patterns of malarial transmission in French Guiana (South America). *Ann Trop Med Parasitol*. 2011;105:107–22.
37. Prussing C, Saavedra MP, Bickersmith SA, Alava F, Guzmán M, Manrique E, et al. Malaria vector species in Amazonian Peru co-occur in larval habitats but have distinct larval microbial communities. *PLoS Negl Trop Dis*. 2019;13:e0007412.
38. de Barros FS, Honorio NA. Deforestation and malaria on the Amazon frontier: larval clustering of *Anopheles darlingi* (Diptera: Culicidae) determines focal distribution of malaria. *Am J Trop Med Hyg*. 2015;93:939–53.
39. Chaves LSM, Berço ES, Conn JE, Laporta GZ, Prist PR, Sallum MAM. Anthropogenic landscape decreases mosquito biodiversity and drives malaria vector proliferation in the Amazon rainforest. *PLoS ONE*. 2021;16:e0245087.
40. Consoal RA, Lourenco-de-Oliveira R. 1994. Principais mosquitos de importância sanitária no Brasil. Fundação Oswaldo Cruz. Brazil: Editora Fiocruz.
41. Farañ ME, Linthicum KJ. A handbook of the Amazonian species of *Anopheles* (*Nyssorhynchus*) (Diptera: Culicidae). *Mosq Syst*. 1981;13:1–81.
42. Forattini OP. *Entomologia Médica*, vol. 1. São Paulo, Brazil: Faculdade de Higiene e Saúde Pública; 1962.
43. Davidson G. Estimation of the survival-rate of anopheline mosquitoes in nature. *Nature*. 1954;174:792–3.
44. Service MW. Community participation in vector-borne disease control. *Ann Trop Med Parasitol*. 1993;87:223–34.
45. de Barros FS, Honorio NA, Aruda ME. Survivorship of *Anopheles darlingi* (Diptera: Culicidae) in relation with malaria incidence in the Brazilian Amazon. *PLoS ONE*. 2011;6:e22388.
46. Bickersmith SA, Lainhart W, Moreno M, Chu VM, Vinetz JM, Conn JE. A sensitive, specific and reproducible real-time PCR method for detection of *Plasmodium vivax* and *P. falciparum* infection in field-collected anophelines. *Mem Inst Oswaldo Cruz*. 2015;110:573–6.
47. Pizzutti F, Pan W, Barbieri A, Miranda JJ, Feingold B, Guedes GR, et al. A validated agent-based model to study the spatial and temporal heterogeneities of malaria incidence in the rainforest environment. *Malar J*. 2015;14:1–19.
48. Venables WN, Ripley BD. *Modern applied statistics with S*. 4th ed. New York: Springer; 2002.
49. Matson R, Rios CT, Chavez CB, Gilman RH, Florin D, Sifuentes VL, et al. Improved molecular technique for the differentiation of neotropical anopheline species. *Am J Trop Med Hyg*. 2008;78:492–8.
50. Rosas-Aguilera A, Gamboa D, Manrique P, Conn JE, Moreno M, Lescano AG, et al. Epidemiology of *Plasmodium vivax* malaria in Peru. *Am J Trop Med Hyg*. 2016;95:133–44.
51. Conn JE, Moreno M, Saavedra M, Bickersmith SA, Knoll E, Fernandez R, et al. Molecular taxonomy of *Anopheles* (*Nyssorhynchus*) *benarrochi* (Diptera: Culicidae) and malaria epidemiology in southern Amazonian Peru. *Am J Trop Med Hyg*. 2013;88:319–24.
52. Prussing C, Emerson KJ, Bickersmith SA, Sallum MAM, Conn JE. Minimal genetic differentiation of the malaria vector *Nyssorhynchus darlingi* associated with forest cover level in Amazonian Brazil. *PLoS ONE*. 2019;14:e0225005.
53. Morales Wieri D, Herrera-Varela M, Albuja M, Quiroga C, Diaz G, Del Aguila Morante C, et al. New records of *Anopheles benarrochi* B (Diptera: Culicidae) in malaria hotspots in the Amazon regions of Ecuador and Peru. *J Med Entomol*. 2021. <https://doi.org/10.1093/jme/tjaa293>.
54. Conn JE, Bickersmith SA, Saavedra MP, Morales JA, Alava F, Diaz Rodriguez GA, et al. Natural infection of *Nyssorhynchus darlingi* and *Nyssorhynchus benarrochi* B with *Plasmodium* during the dry season in the understudied low-transmission setting of Datem del Marañon Province, Amazonian Peru. *Am J Trop Med Hyg*. 2023. <https://doi.org/10.4269/ajtmh.23-0058>.
55. Flores-Mendoza C, Fernandez R, Escobedo-Vargas KS, Vela-Perez Q, Schoeler GB. Natural *Plasmodium* infections in *Anopheles darlingi* and *Anopheles benarrochi* (Diptera: Culicidae) from eastern Peru. *J Med Entomol*. 2004;41:489–94.
56. Monroe A, Moore S, Okumu F, Kiware S, Lobo NF, Koenker H, et al. Methods and indicators for measuring patterns of human exposure to malaria vectors. *Malar J*. 2020;19:207.
57. Manrique P, Miranda-Alban J, Alarcon-Baldeon J, Ramirez R, Carrasco-Escobar G, Herrera H, et al. Microsatellite analysis reveals connectivity among geographically distant transmission zones of *Plasmodium vivax* in the Peruvian Amazon: a critical barrier to regional malaria elimination. *PLoS Negl Trop Dis*. 2019;13:e0007876.
58. Rosas-Aguilera A, Guzman-Guzman M, Chuquiyauro I, Moreno M, Manrique P, Ramirez R, et al. Temporal and micro-spatial heterogeneity in transmission dynamics of co-endemic *Plasmodium vivax* and *Plasmodium falciparum* in two rural cohort populations in the Peruvian Amazon. *J Infect Dis*. 2020;222:1–12.
59. Fontoura PS, Silva MF, da Costa AS, Ribeiro FS, Ferreira MS, Ladela-Andrade S, et al. Monthly biological larviciding associated with a tenfold decrease in larval density in fish farming ponds and reduced community-wide malaria incidence in northwestern Brazil. *Parasit Vectors*. 2021;14:445.
60. Ferreira MJ, Corder RM, Johansen IC, Kattenberg JH, Moreno M, Rosas-Aguilera A, et al. Relative contribution of low-density and asymptomatic infections to *Plasmodium vivax* transmission in the Amazon: pooled analysis of individual participant data from population-based cross-sectional surveys. *Lancet Reg Health Am*. 2022;9:100169.
61. Ramirez RS. Micro-epidemiología molecular des las infecciones que contribuyen a mantener la transmisión de *Plasmodium vivax* en comunidades en vías de eliminación de la región Loreto. San Martín de Porres: Universidad Peruana Cayetano Heredia Facultad de Ciencias y Filosofía; 2024.

62. MINSA, Ministerio de Salud del Perú: Boletín epidemiológico del Perú SE 31-2022, vol. 31. Lima, Perú: Centro Nacional de Epidemiología, Prevención y Control de Enfermedades; 2022.
63. Ministerio de Salud. Documento técnico: plan hacia la malaria en el Perú 2022-2030. 2022:60.
64. Iyer M, Skehon J, de Wildt G, Meza G. A qualitative study on the use of long-lasting insecticidal nets (LLINs) for the prevention of malaria in the Peruvian Amazon. *Malar J*. 2019;18:301.
65. Prado CC, Alvarado-Cabrera LA, Camargo-Ayala PA, Garzón-Ospina D, Camargo M, Soto-De León SC, et al. Behavior and abundance of *Anopheles darlingi* in communities living in the Colombian Amazon riverside. *PLoS ONE*. 2019;14:e0213335.
66. Mmbando AS, Ngowo H, Limwagu A, Kilalangongo M, Rfungo K, Okumu FO. Eave ribbons treated with the spatial repellent, transfluthrin, can effectively protect against indoor-biting and outdoor-biting malaria mosquitoes. *Malar J*. 2018;17:368.
67. Kaindoa EW, Mmbando AS, Shirima R, Hape EE, Okumu FO. Insecticide-treated eave ribbons for malaria vector control in low-income communities. *Malar J*. 2021;20:415.
68. Masaku JP, Pinda M, Killeen GF, Ngowo HS, Pinda PG, Okumu FO. Creating mosquito-free outdoor spaces using transfluthrin-treated chairs and ribbons. *Malar J*. 2020;19:109.
69. Alves FP, Gal LH, Marelli MT, Ribeiro PE, Camargo EP, Da Silva LH. Asymptomatic carriers of *Plasmodium* spp. as infection source for malaria vector mosquitoes in the Brazilian Amazon. *J Med Entomol*. 2005;42:777-9.
70. Lin JT, Saunders DL, Meshnick SR. The role of submicroscopic parasitemia in malaria transmission: what is the evidence? *Trends Parasitol*. 2014;30:183-90.
71. Fouque F, Gabrion P, Carinci R, Issaly J, Girod R. Annual variations in the number of malaria cases related to two different patterns of *Anopheles darlingi* transmission potential in the Maroni area of French Guiana. *Malar J*. 2010;9:80.
72. Oliveira TMP, Laporta GZ, Bergamo ES, Chaves LSM, Antunes JF, Bickersmith SA, et al. Vector role and human biting activity of Anophelinae mosquitoes in different landscapes in the Brazilian Amazon. *Parasit Vectors*. 2021;14:236.
73. Sita DJ, Sanou R, Mitton J, Mwangi EP, Niang A, Sere I, et al. Rapid age-grading and species identification of natural mosquitoes for malaria surveillance. *Nat Commun*. 2022;13:1501.
74. Moreno JE, Rubio-Palis Y, Paez E, Perez E, Sanchez Y. Abundance, biting behaviour and parous rate of anopheline mosquito species in relation to malaria incidence in gold-mining areas of southern Venezuela. *Med Vet Entomol*. 2007;21:339-49.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## II. DISCUSIÓN

### **Influencia de factores espaciales y temporales en la transmisión**

Los hallazgos del presente estudio refuerzan que la transmisión de la malaria en comunidades amazónicas está fuertemente influenciadas por factores espaciales y estacionales que modulan el comportamiento de *Nyssorhynchus darlingi*. La mayor abundancia del vector y la evidencia de transmisión activa estuvo mayormente concentrada en comunidades ribereñas como Lupuna y Santa Emilia. Además, el vector fue predominantemente exofílico, con actividad de picadura en el peridomicilio en las primeras horas de la noche. Se observó también una marcada estacionalidad en la abundancia y la longevidad del vector, especialmente en la temporada de lluvia. Estos resultados confirman que las comunidades ribereñas presentan condiciones ecológicas que favorecen la reproducción sostenida y la transmisión activa del vector. [1,2,3]. Además, refuerzan la necesidad de adaptar las estrategias de intervención vectorial al contexto de cada comunidad, priorizando acciones en comunidades ribereñas, que cuentan con condiciones ecológicas, como criaderos, que facilitan la reproducción del vector [4,5]. Estudios previos también identificaron a *Ny. darlingi* como el principal vector en zonas ribereñas con alta transmisión, como reportaron Rosas-Aguirre et al. y Moreno et al. [6,7]. El uso de indicadores entomológicos como la tasa de inoculación entomológica (EIR), que combina datos de abundancia y hembras infectadas, resulta clave para identificar focos activos de transmisión [8]. En el caso de Santa Emilia, el hallazgo de EIR positivo en múltiples meses justifica la implementación de vigilancia entomológica focalizada y acciones como la distribución de LLINs o IRS, especialmente en la temporada de lluvias, entre diciembre y junio [9,10].

### **Actividad de picadura temprana y comportamiento exofílico**

La concentración de la actividad de picadura del vector en las primeras horas de la noche, entre las 18:00h y las 21:00h, representa un periodo crítico de exposición humana, ya que las personas aún no están protegidas por mosquiteros [11]. Este patrón de picos tempranos ha sido reportado por Girod et al. y Vittor et al. [11,12], quienes destacan la importancia de implementar medidas de control complementarios al uso de LLINs que aborden este periodo de exposición.

El comportamiento exofílico observado en *Ny. darlingi*, particularmente en comunidades ribereñas, también representa un desafío importante para las intervenciones vectoriales tradicionales (LLINs o IRS), que están diseñadas para la protección en el intra-domicilio [2,13]. En el contexto amazónico, las personas suelen estar activas en el peri-domicilio, ya sea por visitas sociales, actividades agrícolas, higiene personal o prácticas religiosas [7]. Estas situaciones de riesgo no están cubiertas adecuadamente por las estrategias actuales de control. Este patrón fue reportado previamente en Loreto (Vittor et al.; Aramburú & Ramal) y también en otras regiones de América Latina, donde se ha reportado la capacidad de *Ny. darlingi* para ajustar su comportamiento en respuesta a condiciones ecológicas o a presiones generadas por las intervenciones [11,14,15].

### **Estacionalidad, longevidad vectorial y riesgo de transmisión**

En El Triunfo, a pesar de presentar baja densidad vectorial, se observó una longevidad máxima de 8.71 días durante marzo (temporada lluviosa), con el 100% de las hembras evaluadas en condición de paridas, este hallazgo sugiere que incluso en comunidades con baja abundancia del vector, puede existir riesgo significativo de transmisión si los mosquitos sobreviven lo suficiente como para completar el

desarrollo esporogónico de *Plasmodium*. Durante la temporada seca, la paridad como la longevidad disminuyeron notablemente, y no se detectaron hembras infectadas con *Plasmodium*, Esto indica una población vectorial envejecida y sostenida, con alto potencial para mantener transmisión. Esta fuerte estacionalidad, resalta la necesidad de reforzar las intervenciones vectoriales de forma anticipada a los meses lluviosos, especialmente en comunidades ribereñas con alta vulnerabilidad ecológica [4,15]. Esta estacionalidad, ya ha sido reportada en estudios previos de Loreto y Alto Amazonas (Vittor et al.; Rosas-Aguirre et al.), y concuerda con evidencia reportada en Brasil y Colombia, donde la abundancia del vector aumenta tras eventos de lluvia e inundación [3,14]. La presencia de hembras infectadas en el peri-domicilio, especialmente en Santa Emilia, refuerza la necesidad de incluir este entorno en los planes de vigilancia y control entomológico [6,16].

Esta situación, que ha sido poco considerada por los programas de control nacional en el Perú, representa una oportunidad para reorientar las intervenciones para lograr una cobertura más integral que considere estos espacios peri-domiciliarios como sitios de alta transmisión. La experiencia en comunidades amazónicas de Brasil y Colombia, que han reportado un patrón similar, demuestran que es factible la adaptación de las estrategias a un entorno peri-domiciliario, mediante el uso de estrategias innovadoras como las estaciones de azúcares tóxicos (ABTS), las barreras físicas, repelentes espaciales y tratamientos focalizados en estructuras externas como letrinas y corrales [17,18]. Las altas tasas de supervivencia y paridad en Santa Emilia y El Triunfo durante la creciente reflejan poblaciones de mosquitos capaces de sostener transmisión [5,19]. Esto sugiere que la vigilancia y las

intervenciones deben ser planificadas con enfoque estacional, reforzando las intervenciones comunitarias antes del incremento esperado de abundancia del vector. Además, se necesita integrar datos de longevidad y paridad como indicadores entomológicos operativos. Incluir estas métricas ayudaría a priorizar comunidades con vectores longevos o con altas tasas de paridad [18].

### **Uso de indicadores operativos y limitaciones logísticas**

La aplicación de estas estrategias en el Perú enfrenta múltiples desafíos. En primer lugar, la limitada cobertura de servicios básicos, dificultades logísticas y acceso restringido a zonas ribereñas [20,16]. Esto obliga a diseñar estrategias logísticamente viables y adaptadas al contexto local. Por ejemplo, la baja detección de hembras infectadas en Lupuna, en meses diferente a abril, podría deberse a subestimaciones de infecciones submicroscópicas, subestimadas por métodos convencionales [6,21].

En este sentido, una recomendación clave sería fortalecer la vigilancia longitudinal mediante el uso de técnicas de detección molecular de hembras infectadas incluso en periodos con baja transmisión [22]. Además, se debe considerar el uso de intervenciones complementarias como el control larvario, especialmente en comunidades donde el acceso a LLINs e IRS es limitado o su adherencia es baja [17].

Los resultados de este estudio también resaltan la necesidad de incorporar indicadores entomológicos como el EIR, que combina la abundancia de picaduras (HBR) y la proporción de hembras infectadas (IR), para orientar de forma más precisa las acciones de control. Actualmente, el Ministerio de Salud del Perú y la GERESA Loreto emplean principalmente el HBR para la planificación de las

intervenciones, pero este solo indica cuántos mosquitos están picando, sin mostrar cuántos están infectados y posiblemente transmitiendo malaria. La evidencia de este estudio sugiere que el EIR permitiría identificar los focos de transmisión y priorizar las intervenciones en zonas de mayor riesgo. Sin embargo, su aplicación enfrenta una limitante importante, para estimar el IR se requieren aplicar pruebas moleculares y/o inmunológicas como ELISA, cuyo costo elevado podría dificultar su implementación rutinaria en el sistema de salud.

### **Impacto de las intervenciones convencionales (LLINs e IRS) sobre el comportamiento vectorial**

Evidencia reciente (2024–2025) indica que a pesar de la alta cobertura de LLINs/ITNs y la aplicación sostenida de IRS, persiste exposición vinculadas a la actividad humana en exteriores durante las primeras horas de la noche y al comportamiento exofílico de los vectores, lo que limita el impacto de las intervenciones centradas solo en interiores [23-25]. En distintos escenarios africanos se ha observado que el IRS puede reducir el riesgo de infección sin generar disminuciones consistentes en la abundancia de mosquitos, y que la transmisión residual continúa asociada a picaduras en exteriores [26,27]. Paralelamente, evaluaciones experimentales han documentado respuestas conductuales de *Anopheles* frente a distintos tipos y estados de ITNs [23]. En la Amazonía peruana, estudios recientes en zonas peri-Iquitos y ensayos en huts experimentales (pequeñas construcciones diseñadas para evaluar el ingreso y el descanso de mosquitos en condiciones controladas) con *Ny. darlingi* aportan información clave sobre sus patrones de entrada y descanso en estructuras tratadas [28], útil para optimizar la focalización temporal y espacial de LLINs/IRS [29,30].

### **Factores socioculturales y necesidad de nuevas herramientas**

También es necesario explorar por qué comunidades como Santa Emilia mantienen un EIR elevado, mientras otras con alta abundancia vectorial, como Lupuna, no, pese a tener alta abundancia vectorial [31]. Esta diferencia podría deberse a factores sociales, culturales o ecológicos que aún no se comprenden del todo y requieren investigación. Otro de los grandes retos en la región amazónica es que las principales estrategias de control están diseñadas para proteger a las personas durante el descanso nocturno [32].

Abordar esta problemática requiere rediseñar las estrategias educativas y logísticas, priorizando la protección personal durante las primeras horas de la noche mediante el uso de repelentes, espirales o barreras físicas en espacios abiertos [33]. Como línea de investigación futura, se plantea evaluar la adherencia y uso real de LLINs en función del horario de descanso, así como explorar nuevas herramientas de intervención aplicables en horarios temprano y en espacios fuera del dormitorio [34].

### **Comportamiento humano, exposición y líneas futuras de investigación**

Es fundamental integrar estudios sobre comportamiento humano y patrones de actividad nocturna para comprender con mayor precisión los momentos y contextos de exposición efectiva al vector [35]. También se debe investigar si este comportamiento de picadura en el peri-domicilio se mantiene en el tiempo o si representa una respuesta adaptativa temporal a intervenciones con uso masivo de LLINs. Estudios observacionales sobre el comportamiento humano en exteriores, podrían guiar intervenciones más eficaces para reducir el contacto humano-vector en contextos donde las herramientas actuales no tienen cobertura [36].

Finalmente, se plantea como hipótesis de investigación futura explorar que factores ecológicos y/o socioculturales modulan la longevidad de los vectores. Dado que una alta paridad indica contacto frecuente con personas e ineficacia de las intervenciones actuales [6]. La variabilidad en el número de hembras disectadas fue una limitante, porque dificulto calcular consistentemente los indicadores de longevidad. Además, el numero bajo de hembras nulíparas en algunos meses limito la estimación de edad del vector. Esta técnica, aunque sencilla, requiere entrenamiento especializado para evitar errores en la identificación del estado ovárico. Para superar esta limitación, se recomienda complementar las identificaciones ováricas con técnicas moleculares y métodos alternativos para estimar la edad vectorial, como la espectrometría infrarroja.

El estudio no incluyo campamentos madereros y/o mineros temporales, como posibles focos adicionales de transmisión. Se sugiere en futuros estudios monitoreo entomológico y parasitológico de estos campamentos, especialmente si existe evidencia de alta movilidad humana entre las comunidades estudiadas.

### III. REFERENCIAS BIBLIOGRÁFICAS

1. Saavedra MP, Conn JE, Álava F, Carrasco-Escobar G, Prussing C, Bickersmith SA, et al. Higher risk of malaria transmission outdoors than indoors by *Nyssorhynchus darlingi* in riverine communities in the Peruvian Amazon. *Parasite Vectors*. 2019; 12:374.
2. Prussing C, Saavedra MP, Bickersmith SA, Álava F, Guzmán M, Manrique E, et al. Malaria vector species in Amazonian Peru co-occur in larval habitats but have distinct larval microbial communities. *PLoS Negl Trop Dis*. 2019;13:e0007412.
3. Bugoro H, Iroofa C, Mackenzie DO, Apairamo A, Hevalao W, Corcoran S, et al. Changes in vector species composition and current vector biology and behavior will favour malaria elimination in Santa Isabel Province, Solomon Islands. *Malar J*. 2011;10:287.
4. Soto-Calle V, Rosas-Aguirre A, Llanos-Cuentas A, Abatih E, DeDeken R, Rodríguez H, et al. Spatio-temporal analysis of malaria incidence in the Peruvian Amazon Region between 2002 and 2013. *Sci Rep*. 2017; 7:40350.
5. Sallum MAM, Conn JE, Bergo ES, Laporta GZ, Chaves LSM, Bickersmith SA, et al. Vector competence, vectorial capacity of *Nyssorhynchus darlingi* and the basic reproduction number of *Plasmodium vivax* in agricultural settlements in the Amazonian Region of Brazil. *Malar J*. 2019;18:117.
6. Rosas-Aguirre A, Guzman-Guzman M, Chuquiyauri R, Moreno M, Manrique P, Ramírez R, et al. Temporal and micro-spatial heterogeneity in transmission dynamics of co-endemic *Plasmodium vivax* and *Plasmodium falciparum* in two rural cohort populations in the Peruvian Amazon. *J Infect Dis*. 2020; jiaa26:1–12.
7. Moreno M, Saavedra MP, Bickersmith SA, Prussing C, Michalski A, Tong Rios C, et al. Intensive trapping of blood-fed *Anopheles darlingi* in Amazonian Peru reveals unexpectedly high proportions of avian blood-meals. *PLoS Negl Trop Dis*. 2017;11:e0005337.
8. Conn JE, Moreno M, Saavedra MP, Bickersmith SA, Prussing C, Gamboa D, et al. Entomological monitoring and impact evaluation of malaria interventions in the Amazon. *Malar J*. 2023;22:117.


9. Ministerio de Salud del Perú (MINSA). Plan Malaria Cero 2017–2021. Lima: MINSA; 2017.
10. Ministerio de Salud del Perú (MINSA). Plan Nacional de Eliminación de la Malaria 2022–2030. Lima: MINSA; 2022.
11. Girod R, Gaborit P, Carinci R, Issaly J, Fouque F. *Anopheles darlingi* bionomics and transmission of *Plasmodium falciparum*, *Plasmodium vivax* and *Plasmodium malariae* in Amerindian villages of the Upper-Maroni Amazonian forest, French Guiana. *Mem Inst Oswaldo Cruz*. 2008;103(7):702–10. doi:10.1590/S0074-02762008000700013.
12. Vittor AY, Pan W, Gilman RH, Tielsch J, Glass G, Shields T, et al. Linking deforestation to malaria in the Amazon: characterization of the breeding habitat of the principal malaria vector, *Anopheles darlingi*. *Am J Trop Med Hyg*. 2006;74(1):3–11.
13. Piedrahita S, Álvarez N, Naranjo-Díaz N, Bickersmith S, Conn J, Correa M. *Anopheles* blood meal sources and entomological indicators related to *Plasmodium* transmission in malaria endemic areas of Colombia. *Acta Trop*. 2022;233:106567.
14. Abreu FVS, Morais MM, Ribeiro SP, Eiras ÁE. Influence of breeding site availability on the spatial distribution of *Anopheles darlingi* in the Brazilian Amazon. *J Vector Ecol*. 2015;40(2):344–55.
15. Carrasco-Escobar G, Gamboa D, Castro MC, Bangdiwala SI, Rodríguez H, Contreras-Mancilla J, et al. Micro-epidemiology and spatial heterogeneity of *P. vivax* parasitaemia in riverine communities of the Peruvian Amazon: a multilevel analysis. *Sci Rep*. 2017; 7:8082.
16. Ramírez RS. Micro-epidemiología molecular de las infecciones que contribuyen a mantener la transmisión de *Plasmodium vivax* en comunidades en vías de eliminación de la región Loreto. Lima: Universidad Peruana Cayetano Heredia; 2024. Tesis de maestría.
17. Sikulu-Lord MT, Maia MF, Milali MP, Mkandawile G, Wirtz RA, Hugo LE, et al. Rapid age-grading and species identification of natural mosquitoes for malaria surveillance. *Sci Rep*. 2018;8:14067.

18. Prado CC, Alvarado-Cabrera LA, Camargo-Ayala PA, Garzón-Ospina D, Camargo M, Soto-De León SC, et al. Behavior and abundance of *Anopheles darlingi* in communities living in the Colombian Amazon riverside. PLoS ONE. 2019;14:e0213335.
19. de Barros FS, Honorio NA, Arruda ME. Survivorship of *Anopheles darlingi* (Diptera: Culicidae) in relation with malaria incidence in the Brazilian Amazon. PLoS ONE. 2011;6:e22388.
20. Parker BS, Paredes Olortegui M, Penataro Yori P, Escobedo K, Florin D, Rengifo Pinedo S, et al. Hyperendemic malaria transmission in areas of occupation-related travel in the Peruvian Amazon. Malar J. 2013;12:178.
21. Lin JT, Saunders DL, Meshnick SR. The role of submicroscopic parasitemia in malaria transmission: what is the evidence? Trends Parasitol. 2014;30(4):183–90.
22. Lainhart W, Bickersmith S, Nadler K, Moreno M, Saavedra M, Chu VM, et al. Evidence for temporal population replacement and the signature of ecological adaptation in a major Neotropical malaria vector in Amazonian Peru. Malar J. 2015; 14:375.
23. Reid E, Govella NJ, Tchouassi DP, Abong'o B, Ouma PO, Killeen GF, et al. Behavioural responses of *Anopheles gambiae* to different insecticide-treated nets in experimental hut trials. Malar J. 2024;23(1):105.
24. Namango IH, Sikulu-Lord MT, Mmbando AS, Mwakalinga SB, Finda MF, Okumu FO. Biting by malaria-infected *Anopheles* and use of interventions during the night in rural Tanzania. PLoS Glob Public Health. 2024;4(3):e0002395.
25. Esayas E, Gowelo S, Assefa M, Vajda EA, Thomsen E, Getachew A, Massebo F, et al. Impact of nighttime human behavior on exposure to malaria vectors and effectiveness of using long-lasting insecticidal nets in the Ethiopian lowlands and highlands. Parasites Vectors. 2024;17(1):520. doi:10.1186/s13071-024-06607-9.
26. Katureebe A, Zinszer K, Arinaitwe E, Rek J, Kakande E, Charland K, Kigozi R, Kilama M, Nankabirwa J, Yeka A, Mawejje H, Mpimbaza A, Katamba H, Donnelly MJ, Rosenthal PJ, Drakeley C, Lindsay SW, Staedke SG, Smith DL,

- Greenhouse B, Kanya MR, Dorsey G. Measures of malaria burden after long-lasting insecticidal net distribution and indoor residual spraying at three sites in Uganda: a prospective observational study. *PLoS Med.* 2016;13(11):e1002167. doi:10.1371/journal.pmed.1002167. PMID:27824885.
27. Pretorius E, Jones RT, Hutchins H, Haines A, Logan JG, Last A, et al. A survey of indoor and outdoor *Anopheles* density, species composition and circumsporozoite rate of malaria vectors on the Bijagós Archipelago, Guinea-Bissau. *Malar J.* 2025;24:195. doi:10.1186/s12936-025-05372-z.
  28. Castro-Llanos F, Stoops CA, Grieco JP, Escobedo-Vargas K, Lopez-Sifuentes V, Lenhart A, Stell FM, et al. Experimental hut resting and entrance behaviour of *Anopheles* (*Nyssorhynchus*) *darlingi* from Zungarococha, Loreto, Northern Peruvian Amazon. *Malar J.* 2025;24(1):267. doi:10.1186/s12936-025-05442-2
  29. Zembere K, Njoroge MM, Kahindi SC, McCann RS, Mukabana WR, Knols BGJ, et al. The potential for attractive toxic sugar baits to complement core malaria intervention strategies: the need for more evidence. *Malar J.* 2024;23:161. doi:10.1186/s12936-024-05161-0.
  30. Ooko M, Omondi CJ, Orondo PW, Kiptoo M, Andagalu B, Ochwedo KO, et al. Malaria burden and residual transmission: two thirds of mosquito bites may not be preventable with current vector control tools on Bioko Island, Equatorial Guinea. *Int J Infect Dis.* 2024;147:107197. doi:10.1016/j.ijid.2024.10.031.
  31. Manrique P, Miranda-Alban J, Alarcon-Baldeon J, Ramirez R, Carrasco-Escobar G, Herrera H, et al. Microsatellite analysis reveals connectivity among geographically distant transmission zones of *Plasmodium vivax* in the Peruvian Amazon: a critical barrier to regional malaria elimination. *PLoS Negl Trop Dis.* 2019;13:e0007876.
  32. Iyer M, Skelton J, de Wildt G, Meza G. A qualitative study on the use of long-lasting insecticidal nets (LLINs) for the prevention of malaria in the Peruvian Amazon. *Malar J.* 2019; 18:301.
  33. Mmbando AS, Ngowo HS, Limwagu AJ, Kilalangongono M, Kifungo K, Okumu FO. Eave ribbons treated with transfluthrin can protect households from malaria vectors in rural Tanzania. *Malar J.* 2018; 17:368.

34. Kaindoa, E.W., Finda, M., Kiplagat, J. *et al.* Housing gaps, mosquitoes and public viewpoints: a mixed methods assessment of relationships between house characteristics, malaria vector biting risk and community perspectives in rural Tanzania. *Malar J* **17**, 298 (2018).
35. de Almeida NCV, Louzada J, Neves M, Carvalho TM, Castro-Alves J, Silvano-Nascimento TF, et al. Larval habitats, species composition and distribution of malaria vectors in regions with autochthonous and imported malaria in Roraima state, Brazil. *Malar J.* 2022; 21:13.
36. Alves FP, Durlacher RR, Menezes MJ, Krieger H, Silva LH, Camargo EP. High prevalence of asymptomatic *Plasmodium vivax* infection in native Amazonian populations. *Am J Trop Med Hyg.* 2002;66(6):641–8.

#### IV. ANEXO (Cuando se trata de un artículo con autor secundario)

 UNIVERSIDAD PERUANA CAYETANO HEREDIA	UNIDAD DE GOBIERNO Y ADMINISTRACIÓN	
	SISTEMA DE ASEGURAMIENTO DE LA CALIDAD	V.01.04 / 27.03.2023
	MANUAL OPERATIVO DE LAS NORMAS Y PROCEDIMIENTOS DEL TRABAJO DE INVESTIGACIÓN PARA LA OBTENCIÓN DE LOS GRADOS ACADÉMICOS DE MAESTRÍA Y DOCTORADO	ESCUELA DE POSGRADO VÍCTOR ALZAMORA CASTRO Página 1 de 30

##### Anexo 8. Formato para validar el desarrollo temático

###### a. Cuando el estudiante no es el primer autor del artículo.

Documento de respaldo del investigador principal

Estimado a quien corresponda

Me dirijo a usted en calidad de **Investigador Científico I de Griffin Laboratory-Wadsworth Center- New York State Department of Health**, con respecto al artículo de investigación titulado *“Effect of spatiotemporal variables on abundance, biting activity and parity of Nyssorhynchus darlingi (Diptera: Culicidae) in peri-Iquitos, Peru”*, de la autoría del estudiante **Marlon Pierino Saavedra Romero**.

Deseo expresar mi respaldo de la información presentada en la declaración jurada del estudiante con respecto al **desarrollo temático del artículo de investigación**. Confirmando que el estudiante es el autor secundario del artículo de investigación y su participación consistió en: **supervisión y conducción del estudio en campo**, así como el apoyo en el análisis de los datos, lo cual se encuentra claramente especificado en la sección **"Author contributions"** del artículo publicado.

Además, puedo afirmar que los datos e información incluidos en el artículo de investigación adjunto se basan en una metodología rigurosa y una revisión crítica de la literatura relevante. Asimismo, puedo atestiguar que el estudiante ha cumplido con todas las normas éticas y de integridad académica en la realización de este artículo, y que la información presentada es precisa y confiable.

Agradezco la oportunidad de respaldar la información presentada en la declaración jurada del estudiante, y me pongo a su disposición para cualquier consulta adicional que pueda tener.

Atentamente,



Sara A. Bickersmith  
Investigador Científico  
Wadsworth Center,  
New York State Department of Health  
Email: sara.bickersmith@health.ny.gov